










Revised checklist of endemic vascular plants of Kazakhstan

Serik A. Kubentayev¹, Daniyar T. Alibekov¹, Yuri V. Perezhogin², Georgy A. Lazkov^{3,4},
Andrey N. Kupriyanov⁵, Alexander L. Ebel^{6,7}, Klara S. Izbastina^{1,8}, Olga V. Borodulina²,
Balsulu B. Kubentayeva¹

1 Astana Botanical Garden, 16 Orynbor Str., 010016, Astana, Kazakhstan

2 Kostanay Regional University named after A. Baitursynova, 47 Baytursynov Str., 110000, Kostanay, Kazakhstan

3 Institute of Biology of the National Academy of Sciences of the Republic of Kyrgyzstan, Bishkek 720011, Kyrgyzstan

4 Research Centre for Ecology and Environment of Central Asia, Bishkek 720040, Kyrgyzstan

5 Federal Research Center of Coal and Coal Chemistry of Siberian Branch of the Russian Academy of Sciences, 18 Sovetsky Ave., 650000, Kemerovo, Russia

6 Tomsk State University, Lenin Ave. 36, 634050 Tomsk, Russia

7 Central Siberian Botanical Garden, Siberian Branch of the Russian Academy of Sciences, Zolotodolinskaya Str. 101, 630090 Novosibirsk, Russia

8 S. Seifullin Kazakh Agrotechnical Research University, 62 Zhengis Ave, 010000, Astana, Kazakhstan

Corresponding author: Balsulu B. Kubentayeva (balsulu1992@mail.ru)

Abstract

We compiled a checklist of endemic vascular plants occurring in Kazakhstan, employing an exhaustive examination of literature sources, herbarium collections, databases and field observations. Our study reveals that 451 taxa can be considered endemic to Kazakhstan, constituting 7.97% of the total vascular plant diversity in the country. These endemic taxa, originating from 139 genera and 34 families, predominantly thrive in the southern regions of Kazakhstan, specifically in the mountain ridges of the Kazakh part of the Tian Shan, including Karatau (123 taxa), Dzungarian Alatau (80 taxa) and Trans-Ili and Kungey Alatau (50 taxa). Notably, 107 endemic species are granted legal protection. Detailed information regarding life form, life cycle, conservation status and geographical distribution across floristic regions was meticulously compiled for each endemic taxon. Of the six groups of life forms, herbs include the highest part of endemic taxa (367 taxa), followed by dwarf semishrubs (25 taxa), shrubs (23 taxa), subshrubs (20 taxa), undershrubs (13 taxa) and trees (3 taxa). The observed life cycles are perennials (408 taxa), annuals (33 taxa) and biennials (10 taxa). This paper serves as a fundamental groundwork for prospective investigations aimed at assessing population sizes and hotspots of plant endemism throughout Kazakhstan, crucial for determining conservation status of endemic plants.

Key words: Biodiversity hotspots, Central Asia, conservation, endemism, floristic division

Introduction

Endemic plants hold particular importance in studying the history of flora and vegetation of diverse geographical regions, since they represent an important element of biodiversity and serve as vital benchmarks for identifying areas of high biodiversity value. In recent years, much attention has been paid to the study of endemic plants, as evidenced by a large number of scientific



Academic editor: Alexander Sennikov

Received: 20 October 2023

Accepted: 5 February 2024

Published: 28 February 2024

Citation: Kubentayev SA, Alibekov DT, Perezhogin YV, Lazkov GA, Kupriyanov AN, Ebel AL, Izbastina KS, Borodulina OV, Kubentayeva BB (2024) Revised checklist of endemic vascular plants of Kazakhstan. *PhytoKeys* 238: 241–279. <https://doi.org/10.3897/phytokeys.238.114475>

Copyright: © Serik A. Kubentayev et al.

This is an open access article distributed under terms of the Creative Commons Attribution License ([Attribution 4.0 International](https://creativecommons.org/licenses/by/4.0/) – CC BY 4.0).

publications (Tojibaev et al. 2020a; Baasanmunkh et al. 2022; Erst et al. 2022; Chung et al. 2023; Villaseñor et al. 2023).

Kazakhstan occupies a central position within Eurasia and holds a notable distinction of being the ninth largest country globally, with 2,724,900 km² of land area. The territory of Kazakhstan is characterised by a remarkable ecological heterogeneity (Abdulina 1999; Akzhigitova et al. 2003), marked by prominent zonal boundaries, notably the demarcation between the cold-temperate and temperate regions of Northern Eurasia and the Irano-Turanian warm region with the Mediterranean-like type of climate, the latter encompassing the southern part of Kazakhstan.

The remarkable diversity of natural conditions in Kazakhstan contributes to the exceptional richness of its flora, its notable originality and a significant number of endemic plant species in Kazakhstan. According to the latest inventory, 5,658 vascular plant species, representing 159 families and 1,067 genera, occur in the country (Abdulina 1999).

The investigation of endemic plant species, which represent a vital and highly vulnerable component of biodiversity, has garnered significant attention in numerous countries. The number of endemic plants in the countries neighbouring Kazakhstan varies, with China exhibiting the highest number of endemic species at 14,939 (Huang et al. 2011). There are over 2,700 endemic taxa in Russia (Kamelin and Budantsev 2019), Mongolia has 102 taxa (Baasanmunkh et al. 2022), Kyrgyzstan has 393 taxa (Lazkov and Sultanova 2014) and Uzbekistan has 378 taxa (Senikov et al. 2016). Based on a comprehensive review of the “Flora of Kazakhstan” (Pavlov 1956–1966), Goloskokov (1969) counted 760 endemic species from 199 genera and 47 families in Kazakhstan. Otherwise, various sources estimated the presence of 709 to 823 species of endemic plants in Kazakhstan (Pavlov 1956–1966; Bykov 1966; Goloskokov 1969; Baitenov 2001; Gemedjieva et al. 2010).

The scientific literature contains a substantial body of work focused on the investigation of endemic taxa in Kazakhstan. However, these publications predominantly revolve around limited geographical areas, such as specific mountain ranges, floristic regions or administrative divisions (Pavlov 1970; Goloskokov 1979; Baitenov 1982; Anapiev 1996; Bakeev and Atikeeva 2015; Ishmuratova et al. 2015, 2016a, 2016b; Sadyrova et al. 2017; Mukhtubaeva et al. 2017; Kupriyanov 2022). Reports focusing on endemic plant species within certain families and genera have also been published, including studies on Poaceae (Kupriyanov et al. 2018), Apiaceae (Klyuykov and Ukrainskaya 2018), Asteraceae (Kupriyanov 2018), Ranunculaceae (Shchegoleva 2019a), Chenopodiaceae (Osmanali et al. 2019) and *Oxytropis* (Perezhogin et al. 2020).

The available information regarding the composition of endemic plant species in Kazakhstan, as documented in the “Flora of Kazakhstan” (Pavlov 1956–1966) and other related sources (Bykov 1966; Goloskokov 1969), is largely outdated. Since then, numerous species previously classified as endemic have been discovered beyond the borders of Kazakhstan or reduced to synonyms. In addition, in the last 10 years alone, more than 25 species of endemic plants have been described as new to science from the territory of Kazakhstan, for example: six species of *Tulipa* (*T. annae* J.de Groot & Zonn, *T. auliekolica* Perezhogin, *T. dianaeverettiae* J.de Groot & Zonn., *T. turgaica* Perezhogin, *T. salsola* Rukšāns & Zubov, *T. ivasczenkoae* Epiktetov & Belyalov) (Epiktetov and Belyalov 2013; Perezhogin 2013; de Groot and Zonneveld 2020; Rukšāns and Zubov 2022); five

apomictic species of *Taraxacum* (*T. atrochlorinum* Kirschner & Štěpánek, *T. corvinum* Kirschner & Štěpánek, *T. dzhungaricola* Kirschner & Štěpánek, *T. sublilacinum* Kirschner & Štěpánek) (Kirschner and Štěpánek 2017); three species of *Allium*: *A. kokuense* R.M.Fritsch, N.Friesen & S.V.Smirn., *A. lepticum* R.M.Fritsch, N.Friesen & S.V.Smirn. and *A. toksanbaicum* N.Friesen & Veselova (Friesen et al. 2021a, 2021b); two species of *Hedysarum* (*H. tarbagataicum* Knjaz. and *H. ulutavicum* Knjaz.) (Knyazev 2019); *Myosotis kazakhstanica* O.D.Nikif. (Nikiforova 2018); *Gagea almaatensis* Levichev, A.Peterson & J.Peterson (Peterson et al. 2016); *Galatella bectauatensis* Kupr. & Koroljuk (Kupriyanov and Korolyuk 2013); *Rhapon-ticoides zaissanica* Kupr., A.L.Ebel & Khrustaleva (Kupriyanov 2020); *Astragalus saphronovae* Kulikov (Kulikov 2014); *Phlomis mindshelkensis* Lazkov (Lazkov 2014); *Phlomoides boroldaica* A.L.Ebel (Ebel et al. 2019); *Fritillaria kolbintsevii* Rukšāns & Zubov (Rukšāns and Zubov 2021); *Galium zaisanicum* Pinzhenina & Kupr. (Pinzhenina and Kupriyanov 2023); *Prangos multicostata* Kljuykov & Lyskov (Lyskov et al. 2016), *Sphaenolobium korovinii* Pimenov & Kljuykov (Pimenov and Kljuykov 2014) and *Nitraria iliensis* Banaev & Tomoshevich (Banaev et al. 2023).

Consequently, the current knowledge regarding the species diversity of endemic plants in Kazakhstan remains poorly available. In order to address this knowledge gap, our research endeavour aimed to compile the checklist of endemic vascular plants in Kazakhstan, based on an extensive analysis of literary sources, comprehensive revision of herbarium collections and data from field observations.

Materials and methods

For the compilation of an endemic plant checklist in Kazakhstan, extensive literature sources were consulted. Initially, nine volumes of the “Flora of Kazakhstan” (Pavlov 1956, 1958, 1960, 1961a, 1961b, 1963, 1964, 1965, 1966) were utilised, alongside the complete list of the country’s flora (Abdulina 1999). The broad-scale inventory of Central Asian plants, “Conspectus Florae Asiae Mediae”, spanning 11 volumes (Kovalevskaya 1968, 1971; Bondarenko and Nabiev 1972; Pakhomova 1974, 1976; Kamelin et al. 1981; Adylov 1983, 1987; Nabiev 1986; Adylov and Zuckerwanik 1993; Khassanov 2015), was also referenced. Additionally, the “Plants of Central Asia” series, consisting of 16 volumes (Grubov 1963–2008), was incorporated. Reports detailing endemic plants within specific geographical and administrative regions of Kazakhstan were used (Pavlov 1970; Goloskokov 1979; Baitenov 1982; Anapiev 1996; Bakeev and Atikeeva 2015; Ishmuratova et al. 2015, 2016a, 2016b; Mukhtubaeva et al. 2017; Sadyrova et al. 2017; Kupriyanov 2022). Furthermore, lists highlighting endemic plants within particular species-rich families and genera were considered, such as Ranunculaceae (Shchegoleva 2019a), Apiaceae (Klyuykov and Ukrainskaya 2018), Asteraceae (Kupriyanov 2018), *Achillea* (Kupriyanov and Kulemin 2023), *Oxytropis* (Perezhogin et al. 2020) and Chenopodiaceae (Suchorukow 2007; Osmanali et al. 2019). Lists encompassing endemic plants within broader geographical regions, which include parts of Kazakhstan, were also reviewed (Tolmachev 1974; Pyak et al. 2008; Tojibaev et al. 2020a; Erst et al. 2022). Additionally, we paid attention to the species described from Kazakhstan and new combinations published from 2013 to 2023, subsequent to the publication of the latest flora list by Abdulina (1999).

Following the compilation of a list of endemic taxa, we conducted a comprehensive re-assessment of the distribution of each species by cross-referencing

published floristic records encompassing the administrative and geographical regions of Kazakhstan (Goloskokov 1949; Stepanova 1962; Karmysheva 1973, 1982; Pavlov 1980; Baitenov 1985; Pugachev 1994; Safronova 1996; Kotukhov 2005; Aralbay et al. 2006; Kadenova et al. 2008; Aipeisova 2012, 2013; Ishmurtova et al. 2016a; Kokoreva et al. 2018; Ivashchenko 2020; Kupriyanov 2020; Sitpayeva et al. 2020; Kubentayev et al. 2021; Orazov et al. 2022, 2024; Khasanov et al. 2023; Kulymbet et al. 2023; Osmonali et al. 2023; Sumbembayev et al. 2023). Furthermore, in order to clarify the presence of presumably endemic plants of Kazakhstan in neighbouring countries, we consulted floristic records of those territories (Kamelin 1990; Yakovlev 2003; Kulikov 2005; Wu et al. 2008; Ryabini-na and Knyazev 2009; Lazkov and Sultanova 2014; Knyazev 2016; Nowak et al. 2020; Vaganov and Shmakov 2020; Sennikov and Tojibaev 2021; Baasanmunkh et al. 2022), as well as publications documenting the discovery of former Kazakhstan endemics outside their native range (Ho and Fu 1993; Yakovlev 2003; Kurtto et al. 2004; German 2005; Mavrodiev et al. 2005; German 2006a, 2006b; German et al. 2006; Belkin 2009; Sennikov et al. 2011; Soskov 2011; German et al. 2012; German et al. 2013; Byalt and Bubyreva 2014; German 2014; Nobis et al. 2014; Pimenov and Kljuykov 2014; Vesselova 2016; German and Al-Shehbaz 2017; Lazkov and Sennikov 2017a, 2017b; Nobis et al. 2017; Pimenov 2017; Golovanov et al. 2018; Golovanov and Knyazev 2019; Ma and Xu 2019; Shchegoleva et al. 2019b; Zolotukhin and Chkalov 2019; Ovchinnikova 2021; Tojibaev et al. 2022; Sennikov and Lazkov 2023; Vaganov 2023; Juramurodov et al. 2024).

To verify endemic taxa distributions, we employed systematic reports detailing the flora of Kazakhstan and its neighbouring regions (Baitenov 1977; German and Chen 2009; Kljuykov et al. 2018; Smirnov et al. 2018; Nobis et al. 2020; Pimenov 2020; German and Veselova 2022; Sennikov et al. 2023). Additionally, we conducted a thorough examination of specimens housed in various herbaria, including LE, MW, TK, TASH, MHA, SVER, KUZ, ALTB, NS, NSK and MOSP (herbarium acronyms according to Thiers (2023)), as well as the data sourced from the Global Biodiversity Information Facility (GBIF 2023), the International Legume Database and Information Service (ILDIS) (Roskov et al. 2009), BrassiBase: Introduction to a novel database on Brassicaceae evolution. Plant & Cell Physiology (Kiefer et al. 2014), World Plants. Synonymic Checklist and Distribution of the World Flora (Hassler 1994–2024) and the Compositae Working Group (CWG) (2023).

Within the scope of this investigation, we provide a list and an analysis of national endemic vascular plants growing strictly within Kazakhstan (see Appendix 1). This study considers two taxonomic levels of endemic plants: species and subspecies; taxa with a rank lower than subspecies were not considered. Additionally, we present a separate list encompassing sub-endemic taxa (see Suppl. material 1). In this paper, sub-endemics refer to taxa that were formerly considered endemics, but subsequently found in a neighbouring country or countries, based on published literature or herbarium material. In addition, we present a list of former endemics of Kazakhstan reclassified as synonyms of taxa with broader geographical distributions (see Suppl. material 2).

The distribution of each endemic taxon in Kazakhstan is given according to the floristic division of the country (Pavlov 1956). This division partitions Kazakhstan's territory into 29 distinct floristic regions and seven subregions (Fig. 1).

The systematic order and taxonomic position of the families are based on the classification of angiosperms by APG IV (2016). The names of the accept-

ed genera and species are mostly in accordance with Plants of the World Online (POWO 2023), with corrections according to recently-published taxonomic revisions. The authorship of species, genera and families has been critically cross-checked against the information provided in the International Plant Names Index (IPNI 2023).

Results

Based on a rigorous revision of endemic vascular plants in Kazakhstan, a total of 451 taxa have been identified as endemic to the country (Appendix 1), which account for 7.97% of the total number (5,658 species) of vascular plants in Kazakhstan (Abdulina 1999). The endemic taxa recognised in this study belong to 139 genera and 34 families. Notably, Kazakhstan is home to five monotypic endemic genera, i.e. *Karatavia* Pimenov & Lavrova, *Botschantzevia* Nabiev, *Tschulaktavia* Bajtenov ex Pimenov & Kljuykov, *Cancriniella* Tzvelev and *Sauria* Bajtenov. No endemic families are present in the country.

The greatest number of endemic taxa is registered in the following families: Asteraceae Bercht. & J.Presl (111 taxa from 29 genera), Fabaceae Lindl. (81 taxa from 6 genera), Apiaceae Lindl. (27 taxa from 13 genera), Lamiaceae Martinov (27 taxa from 8 genera), Boraginaceae Juss. (23 taxa from 10 genera), Brassicaceae Burnett (23 taxa from 8 genera), Amaryllidaceae J.St.-Hil. (22 taxa from 1 genus), Liliaceae Juss. (18 taxa from 3 genera), Rosaceae Juss. (17 taxa from 7 genera), Poaceae Barnhart (15 taxa from 8 genera) and Amaranthaceae Juss. (14 taxa from 10 genera) (Fig. 2A). The remaining 23 families are represented by one to 10 species each.

Genera with the greatest number of endemic taxa are as follows: *Astragalus* L. with 46 taxa, *Oxytropis* DC. with 22 taxa, *Allium* L. with 21 taxa, *Taraxacum* F.H.Wigg. with 20 taxa, *Jurinea* Cass. with 20 taxa, *Tulipa* L. with 13 taxa, *Lappula* Moench with 11 taxa, *Artemisia* L. with nine taxa, *Zygophyllum* L. with nine taxa and *Phlomoides* Moench with nine taxa. The remaining genera are represented by one to seven taxa (Fig. 2B).

The highest concentration of endemic plants was documented in two floristic regions: Karatau, with 123 taxa and Dzungarian Alatau, with 80 taxa. Additionally, a noteworthy range of 30 to 52 endemic taxa were observed in eight other floristic regions, namely Trans-Ili Kungey Alatau (50 taxa), Betpak-Dala (46 taxa), Western Tian Shan (46 taxa), Balkhash-Alakol (46 taxa), Chu-Ili Range (36 taxa), Eastern Upland (35 taxa), Western Upland (31 taxa) and Altai (27 taxa). On the other hand, a comparatively smaller number of endemic plants (not exceeding 5 taxa) were identified in six floristic regions: Caspian Region (4 taxa), Syrt (4 taxa), Kyzylkum (3 taxa), Mangyshlak (2 taxa), Kokchetav (2 taxa) and Buzachi (1 taxon). In the other two floristic regions (Bukeev, Southern Ustyrt), no endemic plants were found (Fig. 3A, B).

Amongst all endemic plants in Kazakhstan, 107 species, constituting 23.7% of the overall number of endemic taxa, are presently under the state-level legal protection (Baitulin 2014).

Amongst the six groups of life forms accepted by the Flora of Kazakhstan (Pavlov 1956), herbs include the greatest part of endemic taxa (367), followed by dwarf semishrubs (25 taxa), subshrubs (20 taxa), shrubs (23 taxa), undershrubs (13 taxa) and trees (3 taxa). The dominant life cycles are perennials (408 taxa), followed by annuals (33 taxa) and biennials (10 taxa).




Figure 1. Map of the floristic division of Kazakhstan (Pavlov 1956): 1 – Syrt, 2 – Tobol-Ishim, 3 – Irtysh, 4 – Semipalatinsk pine forest, 5 – Kokchetav, 6 – Caspian Region, 6a – Bukeyev, 7 – Aktobe, 7a – Mugojar, 8 – Emba, 9 – Turgay, 10 – Western Upland, 10a – Ulutau, 11 – Eastern Upland, 11a – Karkaraly, 12 – Zaysan, 13 – Northern Ustyrt, 13a – Buzachi, 13b – Mangyshlak, 14 – Aral Region, 15 – Kyzylorda, 16 – Betpak-Dala, 17 – Moiynkum, 18 – Balkhash-Alakol, 19 – Southern Ustyrt, 20 – Kyzylkum, 21 – Turkestan, 22 – Altai, 23 – Tarbagatai, 24 – Dzungarian Alatau, 25 – Trans-Ili Kungey Alatau, 25a – Ketmen-Terskey Alatau, 26 – Chu-Ili Range, 27 – Kyrgyz Alatau, 28 – Karatau, 29 – Western Tian Shan.




Figure 2. The largest families by the number of endemic taxa (A). The largest genera by the number of endemic taxa (B).

According to the results of our research, 341 taxa previously considered endemics are recognised as sub-endemics of Kazakhstan (see Suppl. material 1) because they were found in the neighbouring countries. The largest number of plants previously considered endemic to Kazakhstan was found in China – 152 taxa, Kyrgyzstan – 138 taxa, Uzbekistan – 71 taxa, Russia – 59 taxa, Mongolia – 59 taxa, Tajikistan – 31 taxa, Turkmenistan – 19 taxa. The total of 169 former endemic taxa of Kazakhstan were synonymised to taxa with wider distribution ranges (see Suppl. material 2).

Discussion

Based on our critical evaluation of vascular plants of Kazakhstan, 451 taxa are identified as endemic to the country (Appendix 1). This figure corresponds to




Figure 3. Endemic taxa richness in the floristic regions of Kazakhstan (A). The number of endemic taxa in the floristic regions of Kazakhstan (B).

55–63% of the previously-reported numbers, i.e. 709–823 species (Pavlov 1956–1966; Bykov 1966; Goloskokov 1969; Baitenov 2001; Gemedjieva et al. 2010). The substantial difference in the number of endemic taxa compared to earlier publications is due to their reliance on outdated information solely derived from the nine-volume edition of the Flora of Kazakhstan (Pavlov 1956–1966). Our review reveals that 341 taxa previously considered endemics are to be treated as sub-endemics of Kazakhstan (see Suppl. material 1), whereas 169 former endemic taxa were synonymised to taxa with wider distribution ranges (see Suppl. material 2). For example, two most recent publications removed two endemic taxa from Kazakhstan: *Allium valentinae* Pavlov was found in Kyrgyzstan (Sennikov and Lazkov 2023), whereas the generic status of *Pseudomarrubium* was rejected (Zhao et al. 2023). Such examples provide evidence of ongoing taxonomic and floristic studies that are constantly shaping the list of endemic plants of Kazakhstan.

Despite the extensive territory and diverse natural conditions of Kazakhstan, the occurrence of endemic taxa appears highly uneven in the country. The largest number of endemic plants is concentrated in mountainous areas, specifically in the southern and south-eastern parts of Kazakhstan, within the following floristic regions: Karatau (123 taxa), followed by the Dzungarian Alatau (80 taxa) and Trans-Ili Kungey Alatau (50 taxa). These results fully align with the analysis presented by Gemedjieva et al. (2010), who examined the distributions of endemic plants in Kazakhstan, despite their use of outdated taxonomic and distributional data. The distribution pattern of endemic taxa across the territory of Kazakhstan supports the assertion made by Körner (2002) that mountain systems serve as biodiversity and endemism hotspots due to the compression of distinct climatic zones over varying altitudes. According to the latest global analysis of seed plant endemism, the territory of Kazakhstan is assigned to the centre of neo-endemism (Cai et al. 2023). The mountainous regions of Tian Shan in the south and the Altai Mountains in eastern Kazakhstan appear to have favoured this, because the mountainous regions exhibit a great diversity in their plant lineages and, therefore, contribute to high levels of neo-endemism.

The distribution of endemic plants in Kazakhstan is presumably influenced by the geological and climatic history of the territory. Past climate change and geological history help to explain how diversification and relictualisation shape the distribution of neo- and paleoendemism and simultaneously phylogenetic endemism worldwide (Cai et al. 2023). The influence of geological history on speciation, differentiation, migration and extinction of species has been highlighted by many scientists (Takhtajan 1969; Raven and Axelrod 1974; Latham and Ricklefs 1993; Axelrod et al. 1996; Huang et al. 2011). A prime example illustrating this phenomenon is the endemic taxa richness in the ancient Karatau Mountains (123 taxa), located at the westernmost limit of the Tian Shan. The Karatau Mountains possess a complex geological composition, characterised by the presence of the oldest Precambrian shale formations in Central Asia, as well as the Lower Paleozoic formations consisting of metamorphosed limestones and shales, overlain by a quartzite stratum (Kamelin 1990). An important factor in the distribution of endemic plants is long-term climatic stability. Central Asia is known for the presence of ancient plant lineages which survived there due to the continuous history of suitable climatic conditions (e.g. in *Lactuca* s.l.: Kilian et al. (2017)). The influence of geological and climatic history on the distribution of endemic plants in Kazakhstan requires additional research, given the poorly-studied bedrock types in Central Asia.

In the flora of Kazakhstan, herbs include most of the endemic taxa. The life forms of plants reflect their adaptability to environmental conditions and form the units of ecological classification, grouping plants with similar adaptive structures (Aipeisova 2009). According to Yurtsev (1976) and Rabotnov (1978), studies of life forms contribute to the understanding of species biology and their roles within ecosystems. The diversity of life forms represents a cumulative effect of long-term evolutionary processes responding to gradual changes in regional ecological conditions (Keller 1938; Shennikov 1950; Serebryakov 1964).

Amongst endemic plants of Kazakhstan, perennials (408 taxa) are most numerous. A global analysis of the distribution of plant life cycles around the world has shown that annual plants predominate in hot and arid conditions, especially during the long dry season (Poppenwimer et al. 2022). The number of annuals and biennials in Kazakhstan is 43 taxa or 9% of the total number

of endemic plants of Kazakhstan. The distribution of endemic annuals over the territory of Kazakhstan is relatively uniform. The greatest number of annuals is noted in Western Upland (7 taxa), Zaysan (6 taxa) and Karatau (6 taxa). Amongst the families, the greatest number of annual endemic taxa is registered in Boraginaceae Juss. (17 taxa), Brassicaceae Burnett (8 taxa) and Amaranthaceae Juss. (6 taxa); in other families, 1–3 taxa are registered.

Given that numerous endemic plant species have restricted distributions, which makes them more prone to extinction (Myers et al. 2000; Pitman and Jørgensen 2002), it is crucial to emphasise the assessment and protection of such species (Baasanmunkh et al. 2022). Amongst 451 endemic taxa of Kazakhstan, 107 species are currently under state protection. We consider it necessary to further re-assess the status of protection of endemic plants of Kazakhstan using IUCN criteria.

During the critical examination of endemic plants in Kazakhstan, we found that some endemic plants were inaccurately attributed to other countries in the Plants of the World Online (POWO 2023). For instance, *Arthropytum subulifolium* Schrenk and *Atriplex iljinii* Aellen, according to POWO, are supposedly present in Turkmenistan. However, *Arthropytum subulifolium* Schrenk exclusively grows in a narrow region of the Chu-Ili Range in Kazakhstan (Osmanali et al. 2019), while *Atriplex iljinii* is solely found in the Mangistau, Aktobe and Kyzylorda Regions of Kazakhstan (Suchorukow 2007). *Stipa argillosa* Kotukhov and *Thalictrum bykovii* Kotukhov, along with *Gagea azutavica* Kotukhov, are incorrectly recorded in POWO as species native to the Altai Republic in the Russian Federation, whereas these species were described from East Kazakhstan (Kotukhov 1989, 1990, 1998).

At the same time, in the POWO database, certain species were erroneously assigned to Kazakhstan. For instance, the distributions of *Kamelinia tianschanica* F.O.Khass. & I.I. Malzev, *Cousinia xanthiocephala* Tscherneva, *Vicoa krascheninnikovii* Kamelin, *Phlomoides tschimganica* (Vved.) Adylov, Kamelin & Makhm. and *Erysimum aksaricum* Pavlov are limited to Uzbekistan (Kamelin 1976; Kupriyanov 2018; Tojibaev et al. 2020b). *Cousinia balchanica* Tscherneva and *Minuartia palyzanica* Proskur. were described from Turkmenistan (Proskuryakov 1987; Cherneva 1996), while *Taraxacum darschajense* Orazova and *Hedysarum ovczinnikovii* Karimova ex Kovalevsk. were reported to occur in Tajikistan (Orazova 1982; Adylov 1983) and *Crucianella schischkinii* Lincz. was found in both Uzbekistan and Tajikistan (Kamelin 2017). These errors are associated with the difficulty of matching administrative boundaries with plant distribution areas in complicated mountainous territories.

Conclusions

This checklist includes all strictly endemic plants of Kazakhstan, consisting of 451 taxa (species or subspecies) belonging to 139 genera and 34 families. The largest number of endemic taxa is concentrated in mountainous areas, specifically in the southern and south-eastern parts of Kazakhstan.

This paper serves as a fundamental groundwork for prospective investigations aimed at assessing population sizes and numbers of endemic taxa throughout Kazakhstan, crucial for determining their conservation status. Of course, this checklist of plant endemics of Kazakhstan is not final and will be revised in the future as a result of ongoing taxonomic and floristic studies.

Acknowledgements

We thank D.A. German (Altai State University, Russia) for cross-checking the family Brassicaceae and N. Friesen (Osnabrueck University, Germany) for checking the genus *Allium*. We also thank Alexander Sennikov, Arkadiusz Nowak, Denis Melnikov and anonymous reviewers for valuable comments and recommendations. We want to dedicate this article to Kotukhov YA (Altai Botanical Garden, Ridder, Kazakhstan), who made a great contribution to the study of the flora of Kazakhstan.

Additional information

Conflict of interest

All authors declare that they have no competing interests and personal relationships and agree on the contents of the paper.

Ethical statement

No ethical statement was reported.

Funding

This research has been funded by the Science Committee of the Ministry of Science and Higher Education of the Republic of Kazakhstan (Grant No. AP14870712).

Author contributions

Conceptualisation – SAK; methodology – SAK, YVP, DTA; formal analysis – OVB, BBK; writing – preparation of the initial draft – SAK, DTA, VPY, BBK; editing, SAK, DTA, GAL, ALE, ANK; author's supervision – SAK; project administration – SAK, KSI; acquisition of funding – IKS. All authors have read and agreed with the published version of the manuscript.

Author ORCIDs

Serik A. Kubentayev  <https://orcid.org/0000-0002-0369-0591>

Daniyar T. Alibekov  <https://orcid.org/0000-0003-1555-1430>

Yuri V. Perezhogin  <https://orcid.org/0000-0001-6997-8347>

Georgy A. Lazkov  <https://orcid.org/0000-0002-3531-8524>

Andrey N. Kupriyanov  <https://orcid.org/0000-0001-5602-2012>

Alexander L. Ebel  <https://orcid.org/0000-0002-7889-4580>

Klara S. Izbastina  <https://orcid.org/0000-0002-6418-1950>

Olga V. Borodulina  <https://orcid.org/0000-0002-6080-716X>

Balsulu B. Kubentayeva  <https://orcid.org/0000-0002-0841-6516>

Data availability

All of the data that support the findings of this study are available in the main text or Supplementary Information.

References

- Abdulina SA (1999) List of Vascular Plants of Kazakhstan. Almaty, 187 pp. [In Russian]
Adylov TA [Ed.] (1983) Manual of vascular plants of Central Asia. Critical synopsis of the flora of Central Asia, vol. 7. Science Publishers, Uzbek Academy of Sciences, Tashkent, 416 pp. [In Russian]

- Adylov TA [Ed.] (1987) Manual of vascular plants of Central Asia. Critical synopsis of the flora of Central Asia, vol. 9. Science Publishers, Uzbek Academy of Sciences, Tashkent, 400 pp. [In Russian]
- Adylov TA, Zuckerwanik TI [Eds] (1993) Manual of vascular plants of Central Asia. Critical synopsis of the flora of Central Asia, vol. 10. Science Publishers, Uzbek Academy of Sciences, Tashkent, 692 pp. [In Russian]
- Aipeisova SA (2009) Analysis of life forms of plants of the Aktobe floristic district. Bulletin of the Orenburg State University 4: 107–111. [In Russian]
- Aipeisova SA (2012) Synopsis of the flora of the Aktobe floristic district. Aktobe State University, Aktobe, 175 pp. [In Russian]
- Aipeisova SA (2013) Flora of the Aktobe Floristic District. Aktobe State University, Aktobe, 312 pp. [In Russian]
- Akzhigitova N, Brekle Z, Volkova E, Winkler G, Vukhrer B, Ogar Ya, Rachkovskaya I, Saffronova I, Khramtsov V, Makulbekova G, Kurochkina L (2003) Botanical geography of Kazakhstan and Middle Asia (desert region). Komarov Botanical Institute, St. Petersburg, 424 pp. [In Russian]
- Anapiev IM (1996) Endemic, relict and rare plant species of Central Kazakhstan and their protection. In: JS Akyibaev (Ed.) Modern problems of ecology of Central Kazakhstan. E.A. Buketov KarSU, Karaganda, 103–107. [In Russian]
- APG IV (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. Botanical Journal of the Linnean Society 181(1): 1–20. <https://doi.org/10.1111/boj.12385>
- Aralbay NK, Kudabaeva GM, Imanbaeva AA, Veselova PV, Danilov MP, Kurmantaeva AA, Shadrina NV, Kasenova BT (2006) State cadastre of plants of the Mangistau region. Synopsis of higher vascular plants. Printing house LLP “Classics”, Aktau, 229 pp. [In Russian]
- Axelrod DI, Al-Shehbaz IA, Raven PH (1996) History of the modern flora of China. In: Zhang AL, Wu SG (Eds) Floristic Characteristics and Diversity of East Asian Plants. China Higher Education Press and Springer-Verlag, Beijing, China and Berlin, Germany, 43–55.
- Banaev EV, Tomoshevich MA, Khozyaykina SA, Erst AA, Erst AS (2023) Integrative taxonomy of *Nitraria* (Nitrariaceae), description of the new enigmatic species and key to all currently known species. Plants 12(3): 593. <https://doi.org/10.3390/plants12030593>
- Baasanmunkh S, Urgamal M, Oyuntsetseg B, Sukhorukov AP, Tsegmed Z, Son DC, Erst A, Oyundelger K, Kechaykin AA, Norris J, Kosachev P, Ma J-S, Chang KS, Choi HJ (2022) Flora of Mongolia: Annotated checklist of native vascular plants. PhytoKeys 192: 63–169. <https://doi.org/10.3897/phytokeys.192.79702>
- Baitenov MS (1977) *Astragalus* of the Tien Shan. Nauka Press, Almaty, 143 pp. [In Russian]
- Baitenov MS (1982) Endemic plants at high-elevation of the Northern Tien-Shan. Botanicheskie Materialy Gerbariya Botanicheskogo Instituta Botaniki Akademii Nauk Kazakhskoi SSR 12: 3–9. [In Russian]
- Baitenov MS (1985) Alpine flora of the Northern Tien Shan. Science Publishers, Alma-Ata, 230 pp. [In Russian]
- Baitenov MS (2001) Flora of Kazakhstan. Generic complex of flora. Science Publishers, Almaty, 280 pp. [In Russian]
- Baitulin IO [Ed.] (2014) The Red Book of Kazakhstan (plants), Vol. 2(1). Art Print XXI, Astana, 452 pp. [In Russian]
- Bakeev RT, Atikeeva SN (2015) Study of endemic species in Karkaraly district of Karagandy region. Bulletin of Karaganda University: Series – Biology – Medicine – Geography 77: 79–84. [In Russian]

- Belkin DL (2009) The genus *Silene* L. (Smolyovka) in Altai mountain country. *Turczaninowia* 12: 5–16. [In Russian]
- Bochkov DA, Seregin AP (2022) Local floras of Russia: records from literature. Version 1.71. Lomonosov Moscow State University. Occurrence dataset. <https://doi.org/10.15468/rxtjt2>
- Bondarenko ON, Nabiev MM [Eds] (1972) Manual of vascular plants of Central Asia. Critical synopsis of the flora of Central Asia, Vol. 3. Science Publishers, Uzbek Academy of Sciences, Tashkent, 268 pp. [In Russian]
- Byalt VV, Bubyreva VA (2014) Type specimens of the taxa described by V.L. Komarov during his student years which are kept in the St. Petersburg State University Herbarium. *Biological Communications* 3(4): 89–107. [In Russian]
- Bykov BA (1966) Areas of some endem of Kazakhstan. *Botanicheskie Materialy Gerbariya Botanicheskogo Instituta Botaniki Akademii Nauk Kazakhskoi SSR* 4: 3–15. [In Russian]
- Cai L, Kreft H, Taylor A, Schrader J, Dawson W, Essl F, van Kleunen M, Pergl J, Pyšek P, Winter M, Weigelt P (2023) Climatic stability and geological history shape global centers of neo- and paleoendemism in seed plants. *Proceedings of the National Academy of Sciences of the United States of America* 120(30): e2300981120. <https://doi.org/10.1073/pnas.2300981120>
- Cherneva OV (1996) New species of the genus *Cousinia* Cass. (Asteraceae) from Central Asia. *Novosti Sistematiki Vysshikh Rastenii* 30: 153–157. [In Russian]
- Christenhusz MJ, Govaerts R, David JC, Hall T, Borland K, Roberts PS, Fay MF (2013) Tip-toe through the tulips-cultural history, molecular phylogenetics and classification of *Tulipa* (Liliaceae). *Botanical Journal of the Linnean Society* 172(3): 280–328. <https://doi.org/10.1111/boj.12061>
- Chung GY, Jang HD, Chang KS, Choi HJ, Kim YS, Kim H-J, Son DC (2023) A checklist of endemic plants on the Korean Peninsula II. *Korean Journal of Plant Taxonomy* 53(2): 79–101. <https://doi.org/10.11110/kjpt.2023.53.2.79>
- Compositae Working Group (CWG) (2023) Global Compositae Database. <http://www.compositae.org/gcd/>
- de Groot JJ, Zonneveld BJM (2020) Two new tulip species from the Altai mountains, Kazakhstan. *International Rock Gardener* 122: 3–16.
- Ebel AL, Kupriyanov AN, Lashchinskiy NN, Khrustaleva IA (2019) Notes on *Pseudere-mostachys sewerzovii* and a new species of *Phlomoïdes* (Lamiaceae) from South Kazakhstan. *Systematic notes on the materials of P.N. Krylov Herbarium of Tomsk State University* 119: 25–35. <https://doi.org/10.17223/20764103.119.3> [In Russian]
- Epiktetov VG, Belyalov OV (2013) New species of *Tulipa* L. (Liliaceae) from Kazakhstan. *Turczaninowia* 3(3): 5–7. <https://doi.org/10.14258/turczaninowia.16.3.1> [In Russian]
- Erst AS, Nikulin AYU, Nikulin VYU, Ebel AL, Zibzeev EV, Sharples MT, Baasanmunkh S, Choi Hyeok JAE, Olonova MV, Pyak AI, Gureyeva II, Erst TV, Kechaykin A, Luferov A, Maltseva SYU, Nobis M, Lian L, Wang W (2022) Distribution analysis, updated checklist, and DNA barcodes of the endemic vascular flora of the Altai mountains, a Siberian biodiversity hotspot. *Systematics and Biodiversity* 20(1): 1–30. <https://doi.org/10.1080/14772000.2022.2049391>
- Everett D (2013) The genus *Tulipa*. *Tulips of the world*. Kew publishing, Kew, 380 pp.
- Friesen N, Smirnov SV, Leweke M, Seregin AP, Fritsch RM (2021a) Taxonomy and phylogenetics of *Allium* section *Decipientia* (Amaryllidaceae): Morphological characters do not reflect the evolutionary history revealed by molecular markers. *Botanical Journal of the Linnean Society* 197(2): 190–228. <https://doi.org/10.1093/botlinnean/boab023>

- Friesen N, Vesselova P, Osmonaly B., Sitpaeva G, Luferov A, Shmakov A (2021b) *Allium toksanbaicum* (Amaryllidaceae), a new species from Southeast Kazakhstan Phytotaxa 494: 251–267. <https://doi.org/10.11646/phytotaxa.494.3.1>
- GBIF (2023) Global Biodiversity Information Facility. <https://www.gbif.org> [Accessed via GBIF.org on 2023-07-14]
- Gemedjieva N, da Silva JAT, Ryabushkina N (2010) Representation of endemics in floristic subprovinces of Kazakhstan. The Asian and Australasian Journal of Plant Science and Biotechnology 4: 56–63.
- German DA (2005) *Erysimum mongolicum*, a new species from Mongolia, and notes on *E. vassilczenkoi* and *E. kotuchovii* (Brassicaceae). Willdenowia 35(2): 305–313. <https://doi.org/10.3372/wi.35.35212>
- German DA (2006a) About the endemic species *Erysimum transiliense* (Cruciferae). Sistematische Zametki po Materialam Gerbarii Imeni P. N. Krylova pri Tomskom Gosudarstvennom Universitete 97: 23–31. [In Russian]
- German DA (2006b) On the unusual narrow-locality endemic *Isatis frutescens* (Cruciferae). In: Shmakov AI (Ed.) Problems of botany in Southern Siberia and Mongolia: Proceedings of the V International Scientific and Practical Conference (Barnaul, 21–23 November 2006). Azbuka Press, Barnaul, 42–44. [In Russian]
- German DA (2014) Taxonomic remarks on some Asian *Lepidium* s.l. (Brassicaceae). Phytotaxa 186(2): 97–105. <https://doi.org/10.11646/phytotaxa.186.2.4>
- German DA, Al-Shehbaz IA (2017) A taxonomic note on *Sterigmostemum* and related genera (Anchonieae, Cruciferae). Novosti Sistematiki Vysshikh Rastenii 48: 78–83. <https://doi.org/10.31111/novitates/2017.48.78>
- German DA, Chen WL (2009) Notes on the family Brassicaceae in China. Journal of Systematics and Evolution 47(3): 202–219. <https://doi.org/10.1111/j.1759-6831.2009.00022.x>
- German DA, Veselova PV (2022) On the taxonomy and species composition of cruciferous plants (Cruciferae) of Kazakhstan. Turczaninowia 25: 12–16. <https://doi.org/10.14258/turczaninowia.25.4.2>
- German DA, Shmakov AI, Zhang XC, Chen WL, Smirnov SV, Xie L, Kamelin RV, Wang J (2006) Some new floristic findings in Xinjiang, China. Zhiwu Fenlei Xuebao 44(5): 598–603. <https://doi.org/10.1360/aps050105>
- German DA, Chen WL, Smirnov SV, Liu B, Kutzev MG, Wang J, Shmakov AI, Kamelin RV (2012) Plant genera and species new to China recently found in northwest Xinjiang. Nordic Journal of Botany 30(1): 61–69. <https://doi.org/10.1111/j.1756-1051.2011.01341.x>
- German DA, Lazkov GA, Tojibaev KSh, Neveraev UA (2013) New data on diversity and geography of the mustard family (Cruciferae) in Kyrgyzstan and Uzbekistan. Botanicheskii Zhurnal 98(9): 1165–1174. [In Russian]
- Goloskokov VP (1949) Flora and Vegetation of the Highlands of Ile Alatau. Academy of Sciences of Kazakhstan, Alma-Ata, 202 pp. [In Russian]
- Goloskokov VP (1969) Features of species endemism in the flora of Kazakhstan. Botanicheskie Materialy Gerbariya Botanicheskogo Instituta Botaniki Akademii Nauk Kazakhskoi SSR 6: 3–12. [In Russian]
- Goloskokov VP (1979) Endemic plants of the Dzhungarian Alatau. Botanicheskie Materialy Gerbariya Botanicheskogo Instituta Botaniki Akademii Nauk Kazakhskoi SSR 11: 9–21. [In Russian]
- Golovanov YM, Knyazev MS (2019) *Goniolimon callicomum* is a new species in the flora of the Urals. Turczaninowia 22: 31–36. [In Russian] <https://doi.org/10.14258/turczaninowia.22.4.4>

- Golovanov Y, Knyazev M, Yamalov S, Korolyuk A, Lebedeva M (2018) Floristic finds on the territory of the Orenburg region and some adjacent territories (based on the materials of the expeditions of 2016–2017). *Turczaninowia* 21: 131–143. [In Russian] <https://doi.org/10.14258/turczaninowia.21.1.12>
- Grubov VI [Ed.] (1963–2008) *Plants of Central Asia*, Vols 1–16. Science Publishers, Moscow and Leningrad. [In Russian]
- Hassler M (1994–2024) *World Plants. Synonymic Checklist and Distribution of the World Flora*. Version 18.3; last update Jan. 3rd, 2024. <https://www.worldplants.de>
- Ho YC, Fu KT (1993) *Astragalus* subgen. *Phaca*. In: Fu KT (Ed.) *Flora Reipublicae Popularis Sinicae*, Vol. 42(1). Science Press, Beijing, 114–168. [In Chinese]
- Huang JH, Chen JH, Ying JS, Ma KP (2011) Features and distribution patterns of Chinese endemic seed plant species. *Journal of Systematics and Evolution* 49(2): 81–94. <https://doi.org/10.1111/j.1759-6831.2011.00119.x>
- IPNI (2023) *International Plant Names Index*. Published on the Internet. The Royal Botanic Gardens Kew, Harvard University Herbaria & Libraries and Australian National Botanic Gardens. <http://www.ipni.org>
- Ishmuratova MYu, Tleukenova SU, Dodonova ASH, Gavrilkova EA (2015) Analysis of endemic plant species of the flora of Karaganda Region. *Actual Problems of the Present* 2(8): 221–225. [In Russian]
- Ishmuratova MYu, Tleukenova SU, Dodonova ASH, Gavrilkova EA (2016a) To the evaluation of the modern species composition of the endemic plants of the flora of the Karaganda region. *Problems of Botany of South Siberia and Mongolia* 15: 381–385. [In Russian]
- Ishmuratova MYu, Myrzaly GZh, Ivlev VI, Matveev AN (2016b) *Flora of Ulytau Mountains (The Central Kazakhstan)*. “Bolashak” Academy, Karagandy, 127 pp. [In Russian]
- Ivashchenko AA (2020) The flora of Sairam-Ugam State National Natural Park (Kazakhstan). *Research of Siberia and Kazakhstan* 26: 52–63. [In Russian]
- Juramurodov IJ, Tojibaev KSh, Sennikov AN, Makhmudjanov DI (2024) The genus *Hedysarum* (Fabaceae; Hedysareae) in Uzbekistan. *Plant Diversity of Central Asia* 3(1): 1–90. https://doi.org/10.54981/PDCA/vol3_iss1/a
- Kadenova AB, Kamkin VA, Erzhanov NT, Kamkina EV (2008) *Flora and Vegetation of the Bayanaul State National Natural Park*. Kereku, Pavlodar, 383 pp. [In Russian]
- Kamelin RV (1976) A new species of the genus *Vicoa* Cass. and other novelties from the Nurata Mountains. *Novosti Sistematiki Vysshikh Rastenii* 13: 244–246. [In Russian]
- Kamelin RV (1990) *Flora of the Syrdarya Karatau*. Science Publishers, Leningrad, 184 pp. [In Russian]
- Kamelin RV (2017) Flora of coloured clays of Middle Asia (brief analysis and problems of genesis). *Turczaninowia* 20: 125–151. [In Russian] <https://doi.org/10.14258/turczaninowia.20.4.14>
- Kamelin RV, Budantsev AL (2019) *Plant World*. Great Russian Encyclopedia. Electronic Version. <https://old.bigenc.ru/biology/text/5554248> [Accessed on 15 May 2023]
- Kamelin RV, Kovalevskaya SS, Nabiev MM [Eds] (1981) *Manual of vascular plants of Central Asia. Critical synopsis of the flora of Central Asia*, Vol. 6. Science Publishers, Uzbek Academy of Sciences, Tashkent, 396 pp. [In Russian]
- Karmysheva NK (1973) *Flora and vegetation of the Aksu-Dzhabagly reserve (Talas Alatau)*. Academy of Sciences of Kazakhstan, Alma-Ata, 173 pp. [In Russian]
- Karmysheva NK (1982) *Flora and vegetation of the western spur of Talass Alatau*. Science Publishers, Alma-Ata, 160 pp. [In Russian]

- Keller BA (1938) Plant and environment. Ecological types and life forms. In: Zinserling YuD (Ed.) Vegetation of the USSR, Vol. 1. Academy of Sciences of the USSR, Moscow and Leningrad, 1–13.
- Khasanov FO, Kubentaev SA, Turdiboev OA (2023) Taxonomical revision and typification of species of *Allium* L. sect. *Multicaulea* and *Unicaulea*. Problems of Botany of South Siberia and Mongolia 22(1): 391–399. <https://doi.org/10.14258/pbssm.2023076>
- Khassanov FO (2015) Conspectus Florae Asiae Mediae, Vol. 11. Fan Publishers, Tashkent, 382 pp. [In Russian]
- Kiefer M, Schmickl R, German DA, Lysak M, Al-Shehbaz IA, Franzke A, Mummenhoff K, Stamatidis A, Koch MA (2014) BrassiBase: Introduction to a novel database on Brassicaceae evolution. Plant & Cell Physiology 55(1): e3. <https://doi.org/10.1093/pcp/pct158>
- Kilian N, Sennikov A, Wang ZH, Gemeinholzer B, Zhang JW (2017) Sub-Paratethyan origin and Middle to Late Miocene principal diversification of the Lactucinae (Compositae: Cichorieae) inferred from molecular phylogenetics, divergence-dating and biogeographic analysis. Taxon 66(3): 675–703. <https://doi.org/10.12705/663.9>
- Kirschner J, Štěpánek J (2017) A revision of *Taraxacum* sect. *Atrata*, a dandelion group centred in the Middle Asia, and the problem of *Taraxacum brevirostre*. Phytotaxa 305(4): 225–261. <https://doi.org/10.11646/phytotaxa.305.4.1>
- Kljuykov EV, Lyskov DF, Ukrainskaja UA (2018) An annotated checklist of the endemic Apiaceae of Kazakhstan. Phytotaxa 360(3): 237–254. <https://doi.org/10.11646/phytotaxa.360.3.4>
- Klyuykov EV, Ukrainskaya UA (2018) Endemic Umbellifers of Kazakhstan. Botanical studies of Siberia and Kazakhstan 24: 3–14. [In Russian]
- Knyazev MS (2016) The lost *Astragalus kustanaicus* (Section Dissitiflori, Fabaceae) found. Botanical Journal. 101: 814–818. <https://doi.org/10.1134/S000681361607005X> [In Russian]
- Knyazev MS (2019) Species of *Hedysarum ucrainicum* affinity (section *Multicaulia*, Fabaceae). Novosti Sistematiki Vysshikh Rastenii 50: 132–140. [In Russian] <https://doi.org/10.31111/novitates/2019.50.132>
- Kokoreva II, Sytpaeva GT, Kudabaeva GM, Otradnykh IG, S'edina IA (2018) Flora of the Northern Tien Shan. The Chu-Ili Range Mountains. Korea National Arboretum, Pocheon, 319 pp. [In Russian]
- Körner C (2002) Mountain biodiversity, its causes and functions: An overview. In: Körner C, Spehn EM (Eds) Mountain biodiversity: a global assessment. Parthenon, London, 3–20. <https://doi.org/10.4324/9780429342585-1>
- Kotukhov YuA (1989) A new species of the genus *Gagea* (Liliaceae) from Southern Altai. Botanicheskii Zhurnal 74: 1663–1964. [In Russian]
- Kotukhov YuA (1990) New species of the genus *Thalictrum* (Ranunculaceae) from Southern Altai. Botanicheskii Zhurnal 75: 416–419. [In Russian]
- Kotukhov YA (1998) New species of grasses (Poaceae) from Southern Altai, Saur and Tarbagatai. Turczaninowia 1: 7–21. [In Russian]
- Kotukhov YuA (2005) List of vascular plants of the Kazakhstan Altai. Botanical Studies of Siberia and Kazakhstan 11: 11–83. [In Russian]
- Kovalevskaya SS [Ed.] (1968) Manual of vascular plants of Central Asia. Critical synopsis of the flora of Central Asia, Vol. 1. Science Publishers, Tashkent, 229 pp. [In Russian]
- Kovalevskaya SS [Ed.] (1971) Manual of vascular plants of Central Asia. Critical synopsis of the flora of Central Asia, Vol. 2. Tashkent, 365 pp. [In Russian]
- Krylov PN (1933) Flora of Western Siberia, 2nd edn., Vol. 7. Tomsk Botanical Department, Russian Society of Naturalists, Tomsk, 387 pp. [In Russian]

- Kubentayev SA, Levichev IG, Sitpayeva GT, Mukhtubayeva SK, Izbastina KS (2021) First records of five species of *Gagea* (Liliaceae), new in the Kazakhstan flora. *Nature Conservation Research* 6(4): 112–114. <https://dx.doi.org/10.24189/ncr.2021.045>
- Kulikov PV (2005) Synopsis of the flora of Chelyabinsk Region (vascular plants) Geotur Publishers, Yekaterinburg and Miass, 537 pp. [In Russian]
- Kulikov PV (2014) A new species of *Astragalus* (Fabaceae) from Plateau Ustyurt (Kazakhstan). *Botanicheskii Zhurnal* 99(9): 1056–1066. [In Russian] <https://doi.org/10.1134/S1234567814090092>
- Kulymbet K, Mukhitdinov N, Kubentayev S, Tynybayeva K, Tastanbekova A, Kurmanbayeva M, Gafforov Y, Kaparbay R, Zhumagul M (2023) The current state of the cenopopulations of *Adonis tianschanica* (Adolf) Lipsch. (Ranunculaceae) in Southeast Kazakhstan. *Biodiversitas* 24: 4359–4372. <https://doi.org/10.13057/biodiv/d240817>
- Kupriyanov AN (2018) Endemism of the flora of Kazakhstan (Asteraceae Bercht. et J.Presl). *Botanical Studies of Siberia and Kazakhstan*. 24: 15–44. [In Russian]
- Kupriyanov AN (2020) Synopsis of the flora of the Kazakh uplands. Academic Publishing House “Geo”, Novosibirsk, 358 pp. [In Russian]
- Kupriyanov AN, Ebel AL, Khrustaleva IA (2020) *Rhaponticoides zaissanica* (Asteraceae) – a new species from Kazakhstan. *Systematic notes on the materials of P.N. Krylov Herbarium of Tomsk State University* 121: 24–31.
- Kupriyanov AN (2022) Endemic plants of the Kazakh Small Hills. *Botanical Studies of Siberia and Kazakhstan* 28: 26–37. [In Russian]
- Kupriyanov AN, Korolyuk EA (2013) *Galatella bectauatense* (Asteraceae), a new species from Central Kazakhstan. *Turczaninowia* 16(3): 15–18. <https://doi.org/10.14258/turczaninowia.16.3.3>
- Kupriyanov AN, Kulemin YuE (2023) Synopsis of the genus *Achillea* L. (Asteraceae) in Kazakhstan. *Turczaninowia* 26(2): 190–201. <https://doi.org/10.14258/turczaninowia.26.2.15> [In Russian]
- Kupriyanov AN, Gudkova PD, Kruchkova EA (2018) Endemism of the Kazakhstan flora: Poaceae Barnhart. *Ukrainian Journal of Ecology* 8(4): 393–405.
- Kurtto A, Lampinen R, Junikka L (2004) *Atlas Florae Europaeae*. Distribution of vascular plants in Europe, Vol. 13. Finnish Museum of Natural History, Helsinki, 320 pp.
- Latham RE, Ricklefs RE (1993) Continental comparisons of temperate-zone tree species diversity. In: Ricklefs, RE Schluter D (Eds) *Species diversity in ecological communities: Historical and geographical perspectives*. University of Chicago Press, Chicago, 294–314.
- Lazkov GA (2014) Taxonomy and distribution of *Phlomis salicifolia* Regel sensu lato and related species of *Phlomis* sect. *Oxyphlomis* (Benth.) Kamelin et Makhm. (Lamiaceae) in Central Asia. — *Novitates Systematicae Plantarum Vascularium* 45: 84–90. <https://doi.org/10.31111/novitates/2014.45.84> [In Russian]
- Lazkov GA, Sennikov AN (2017a) Taxonomy of two blue-flowered juno irises (*Iris* subgen. *Scorpiris*, Iridaceae) from the Western Tian-Shan. *Annales Botanici Fennici* 54(4–6): 297–305. <https://doi.org/10.5735/085.054.0613>
- Lazkov GA, Sennikov AN (2017b) Taxonomic corrections and new records in vascular plants of Kyrgyzstan, 5. *Memoranda Societatis Pro Fauna et Flora Fennica* 93: 79–100.
- Lazkov GA, Sultanova BA (2014) Cadastre of the flora of Kyrgyzstan. Vascular plants. PROON, Bishkek, 125 pp. [In Russian]
- Lyskov DF, Kljuykov EV, Samigullin TH, Ukrainskaja UA, Khrustaleva IA (2016) *Prangos multicostata* (Apiaceae), a new species from eastern Kazakhstan. *Phytotaxa* 277(1): 68–76. <https://doi.org/10.11646/phytotaxa.277.1.6>

- Ma K, Xu Z (2019) NSII (China National Specimen Information Infrastructure, 国家标本资源共享平台). Chinese Academy of Sciences (CAS). Metadata dataset. <https://doi.org/10.15468/kmob80> [Accessed via GBIF.org on 2024-01-04]
- Malyshev LI [Ed.] (1994) Flora of Siberia, Vol. 9. Science Publishers, Novosibirsk, 280 pp. [In Russian]
- Malyshev LI [Ed.] (1997) Flora of Siberia, Vol. 11. Science Publishers, Novosibirsk, 296 pp. [In Russian]
- Mavrodiev EV, Tancig M, Sherwood AM, Gitzendanner MA, Rocca J, Soltis PS, Soltis DE (2005) Phylogeny of *Tragopogon* L. (Asteraceae) based on internal and external transcribed spacer sequence data. International Journal of Plant Sciences 166(1): 117–133. <https://doi.org/10.1086/425206>
- Mukhtubaeva SK, Nelina NV, Sitpayeva GT, Kudabaeva GM, Veselova PV, Bilibaeva BK, Zhumadilova A (2017) Rare, endemic, relict and endangered plant species of Northern Tien Shan (Kungey and Kyrgyz Alatau). Reports of the National Academy of Science, Republic of Kazakhstan 6: 103–110. [In Russian]
- Myers N, Mittermeier RA, Mittermeier CG, da Fonseca GAB, Kent J (2000) Biodiversity hotspots for conservation priorities. Nature 403(6772): 853–858. <https://doi.org/10.1038/35002501>
- Nabiev MM [Ed.] (1986) Manual of vascular plants of Central Asia. Critical synopsis of the flora of Central Asia, Vol. 8. Science Publishers, Uzbek Academy of Sciences, Tashkent, 192 pp. [In Russian]
- Nikiforova O (2018) *Myosotis kazakhstanica* (Boraginaceae) – a new species from Kazakhstan. Turczaninowia 21(3): 89–96. <https://doi.org/10.14258/turczaninowia.21.3.12>
- Nobis M, Nowak A, Nobis A, Paszko B, Piwowarczyk R, Nowak S, Plášek V (2014) Contribution to the flora of Asian and European countries: New national and regional vascular plant records. Acta Botanica Gallica 161(1): 81–89. <https://doi.org/10.1080/12538078.2013.871209>
- Nobis M, Erst A, Nowak A, Shaulo D, Olonova M, Kotukhov Y, Doğru-Koca A, Dönmez AA, Király G, Ebel AL, Kushunina M, Piwowarczyk R, Sukhorukov AP, Nobis A, Verloove F, Zalewska-Gałosz J, Zare G, Burri J, Caković D, Jędrzejczak E, Jogan N, Klichowska E, Pliszko A, Popovich AV, Stešević D, Šilc U, Tupitsyna N, Vasjukov VM, Wang W, Werner P, Wolanin MN, Wolanin MM, Xiang KL (2017) Contribution to the flora of Asian and European countries: New national and regional vascular plant records. Botany Letters 164(1): 23–45. <https://doi.org/10.1080/23818107.2016.1273134>
- Nobis M, Gudkova P, Nowak A, Sawicki J, Nobis A (2020) A synopsis of the genus *Stipa* (Poaceae) in Middle Asia, including a key to species identification, an annotated checklist, and phytogeographic analyses. Annals of the Missouri Botanical Garden 105(1): 1–63. <https://doi.org/10.3417/2019378>
- Nowak A, Nobis M, Nowak S, Nobis A, Wróbel A, Swierszcz S, Klichowska E, Dembicz I, Kusza G (2020) Illustrated flora of Tajikistan and adjacent areas, Vol. 2. Polish Academy of Sciences, Botanical Garden Center for Biological Diversity Conservation and Polish Botanical Society, Warsaw, 768 pp. <https://digitalcollection.zhaw.ch/handle/11475/20859>
- Orazov A, Myrzagaliyeva A, Mukhitdinov N, Tustubayeva S (2022) Callus induction with 6-BAP and IBA as a way to preserve *Prunus ledebouriana* (Rosaceae), and endemic plant of Altai and Tarbagatai, East Kazakhstan. Biodiversitas 23: 3178–3184. <https://doi.org/10.13057/biodiv/d230645>
- Orazov A, Yermagambetova M, Myrzagaliyeva A, Mukhitdinov N, Tustubayeva S, Turuspekov Y, Almerikova S (2024) Plant height variation and genetic diversity between

- Prunus ledebouriana* (Schlecht.) YY Yao and *Prunus tenella* Batsch based on using SSR markers in East Kazakhstan. PeerJ 12: e16735. <https://doi.org/10.7717/peerj.16735>
- Orazova AO (1982) A new species of dandelion from Central Asia. Botanicheskie Materialy Gerbariya Botanicheskogo Instituta Botaniki Akademii Nauk Kazakhskoi SSR 12: 58–59. [In Russian]
- Osmanali BB, Veselova PV, Kudabayeva GM (2019) Features of the distribution of endemic species of the family Chenopodiaceae Vent. in Kazakhstan. Problems of Botany of Southern Siberia and Mongolia 18: 343–346. [In Russian] <https://doi.org/10.14258/pbssm.2019068>
- Osmonali BB, Vesselova PV, Kudabayeva GM, Skaptsov MV, Shmakov AI, Friesen N (2023) Phylogeny and Flow Cytometry of the Genus *Kalidium* Moq. (Amaranthaceae s.l.) in Kazakhstan. Plants. 2023 12(14): 2619. <https://doi.org/10.3390/plants12142619>
- Ovchinnikova SV (2021) Critical notes on kinship group *Lappula subcaespitosa* (Boraginaceae). Flora and Vegetation of Asian Russia 1(1): 3–16. <https://doi.org/10.15372/RMAR20210101> [In Russian]
- Pakhomova MG [Ed.] (1974) Manual of vascular plants of Central Asia. Critical synopsis of the flora of Central Asia, Vol. 4. Science Publishers, Tashkent, 273 pp. [In Russian]
- Pakhomova MG [Ed.] (1976) Manual of vascular plants of Central Asia. Critical synopsis of the flora of Central Asia, Vol. 5. Science Publishers, Uzbek Academy of Sciences, Tashkent, 274 pp. [In Russian]
- Pavlov NV [Ed.] (1956) Flora of Kazakhstan, Vol. 1. Academy of Sciences of the Kazakh SSR, Alma-Ata, 354 pp. [In Russian]
- Pavlov NV [Ed.] (1958) Flora of Kazakhstan, Vol. 2. Academy of Sciences of the Kazakh SSR, Alma-Ata, 292 pp. [In Russian]
- Pavlov NV [Ed.] (1960) Flora of Kazakhstan, Vol. 3. Academy of Sciences of the Kazakh SSR, Alma-Ata, 460 pp. [In Russian]
- Pavlov NV [Ed.] (1961a) Flora of Kazakhstan, Vol. 4. Academy of Sciences of the Kazakh SSR, Alma-Ata, 548 pp. [In Russian]
- Pavlov NV [Ed.] (1961b) Flora of Kazakhstan, Vol. 5. Academy of Sciences of the Kazakh SSR, Alma-Ata, 515 pp. [In Russian]
- Pavlov NV [Ed.] (1963) Flora of Kazakhstan, Vol. 6. Academy of Sciences of the Kazakh SSR, Alma-Ata, 465 pp. [In Russian]
- Pavlov NV [Ed.] (1964) Flora of Kazakhstan, Vol. 7. Academy of Sciences of the Kazakh SSR, Alma-Ata, 497 pp. [In Russian]
- Pavlov NV [Ed.] (1965) Flora of Kazakhstan, Vol. 8. Science Publishers, Kazakh SSR, Alma-Ata, 447 pp. [In Russian]
- Pavlov NV [Ed.] (1966) Flora of Kazakhstan, Vol. 9. Science Publishers, Kazakh SSR, Alma-Ata, 640 pp. [In Russian]
- Pavlov VN (1970) Endemism of the flora of Western Tian-Shan. Botanicheskii Zhurnal (Leningrad) 9: 1232–1241. [In Russian]
- Pavlov VN (1980) Vegetation of Western Tian-Shan. Moscow State University Press, Moscow, 248 pp. [In Russian]
- Perezhogin YuV (2013) New species of tulips from Northern Kazakhstan. Botanicheskii Zhurnal 98: 1558–1563. [In Russian]
- Perezhogin YuV, Yerokhin NG, Tarasov MS, Tanat D (2020) Endemic species of the genus *Oxytropis* in Kazakhstan. Bulletin of KSPI 4: 32–37. [In Russian]
- Peterson A, Harpke D, Levichev IG, Beisenova S, Schnittler M, Peterson J (2016) Morphological and molecular investigations of *Gagea* (Liliaceae) in southeastern Kazakh-

- stan with special reference to putative altitudinal hybrid zones. *Plant Systematics and Evolution* 302(8): 985–1007. <https://doi.org/10.1007/s00606-016-1313-7>
- Pimenov MG (2017) Updated checklist of Chinese Umbelliferae: nomenclature, synonymy, typification, distribution. *Turczaninowia* 20: 106–239. [In Russian] <https://doi.org/10.14258/turczaninowia.20.2.9>
- Pimenov MG (2020) Updated checklist of the Umbelliferae of Middle Asia and Kazakhstan: nomenclature, synonymy, typification, distribution. *Turczaninowia* 23: 127–257. [In Russian] <https://doi.org/10.14258/turczaninowia.23.4.12>
- Pimenov MG, Kljuykov EV (2014) New species of Umbelliferae from Central Asia and Southern Kazakhstan. *Botanicheskii Zhurnal* 99(5): 594–609. [In Russian] <https://doi.org/10.11646/phytotaxa.99.1.2>
- Pinzhenina EA, Kupriyanov AN (2023) *Galium zaisanicum* (Rubiaceae), a new species from Zaisan Depression (Republic of Kazakhstan). *Flora and Vegetation of Asian Russia* 16(1): 72–82. <https://doi.org/10.15372/RMAR20230107> [In Russian]
- Pitman NCA, Jørgensen PM (2002) Estimating the size of the world's threatened flora. *Science* 298(5595): 989–989. <https://doi.org/10.1126/science.298.5595.989>
- Poppenwimer T, Mayrose I, DeMalach N (2022) Revising the global biogeography of plant life cycles. *BioRxiv*. <https://doi.org/10.1101/2022.09.13.507878>
- POWO (2023) Plants of the World Online. Facilitated by the Royal Botanic Gardens, Kew. Published on the Internet. <http://www.plantsoftheworldonline.org> [Retrieved 09 April 2023]
- Proskuryakov GM (1987) New *Minuartia* from the *Sabulina* section. *Byulleten'. Glavnogo Botanicheskogo Sada* 143: 33–35. [In Russian]
- Pugachev PG (1994) Pine forests of the Turgai depression. Kostanay Press, Kostanay, 406 pp. [In Russian]
- Pyak AI, Shaw SC, Ebel AL, Zverev AA, Hodgson JG, Wheeler CD, Gaston KJ, Morenko MO, Revushkin AS, Kotukhov YA, Oyunchimeg D (2008) Endemic plants of the Altai Mountain country. Princeton University Press, 368 pp.
- Rabotnov TA (1978) *Phytocenology*. Moscow State University, Moscow, 384 pp. [In Russian]
- Raven PH, Axelrod DI (1974) Angiosperm biogeography and past continental movements. *Annals of the Missouri Botanical Garden* 61(3): 539–673. <https://doi.org/10.2307/2395021>
- Roskov Y, Bisby FA, Zarucchi JL, Schrire BD, White RJ (2009) ILDIS World Database of Legumes (version 10.01 May 2007) CD-ROM. Species 2000 & ITIS Catalogue of Life: Annual Checklist. Species 2000, Reading.
- Rukšāns J, Zubov D (2021) *Fritillaria kolbintsevii* (subgen. *Fritillaria*; Liliaceae): a new fritillary species from Zhetysu area of south-eastern Kazakhstan. *International Rock Gardener, Scottish Rock Garden Club* 141: 1–45.
- Rukšāns J, Zubov D (2022) Two new *Tulipa* species from sect. *Biflores* (subgen. *Eriostemon*, Liliaceae) described from Iran, Zagros Mountains and Kazakhstan, Zhetysu. *International Rock Gardener* 148: 2–51.
- Ryabinina ZN, Knyazev MS (2009) Key to vascular plants of the Orenburg Region. KMK Scientific Press, Moscow, 758 pp. [In Russian]
- Sadyrova GA, Baizhygitov DK, Dzhamilova SM, Inelova ZA, Yerubayeva GK (2017) Endemics and Subendemics Species Diversity of Ketpen Ridge Flora. *Online Journal of Biological Sciences* 17(4): 299–308. <https://doi.org/10.3844/ojbsci.2017.299.308>
- Safronova IN (1996) Mangyshlak deserts (an outline of their vegetation). *Proceedings of the V.L. Komarov Botanical Institute of the Russian Academy of Sciences* 18: 1–211. [In Russian]

- Sennikov A, Lazkov A (2023) Taxonomic revision of the *Allium filidens* group (Amaryllidaceae) in Kyrgyzstan. *Nordic Journal of Botany* 2023(9): e04050. <https://doi.org/10.1111/njb.04050>
- Sennikov AN, Tojibaev KSh (Eds) (2021) Checklist of vascular plants of the Tian-Shan Mountain System. Korea National Arboretum, Pocheon, 607 pp. [In Russian]
- Sennikov AN, Lazkov GA, Uotila P, Weber HE (2011) Taxonomic corrections and new records in vascular plants of Kyrgyzstan. *Memoranda Societatis Pro Fauna et Flora Fennica* 87: 41–64.
- Sennikov AN, Tojibaev KS, Khassanov FO, Beshko NY (2016) The Flora of Uzbekistan Project. *Phytotaxa* 282(2): 107–118. <https://doi.org/10.11646/phytotaxa.282.2.2>
- Sennikov AN, Khassanov FO, Ortikov E, Kurbonaliyeva M, Tojibaev KS (2023) The genus *Iris* L. s.l. (Iridaceae) in the Mountains of Central Asia biodiversity hotspot. *Plant Diversity of Central Asia* 2(1): 1–104. https://doi.org/10.54981/PDCA/vol2_iss1/a1
- Serebryakov IG (1964) Life forms of plants and their study. In: Lavrenko EM, Korchaagin AA (Eds) *Field Geobotany*, Vol. 3. Science Publishers, Moscow and Leningrad, 146–205. [In Russian]
- Shchegoleva NV (2019a) Endemic species of *Ranunculus* in the flora of Kazakhstan. *Botanical studies of Siberia and Kazakhstan* 25: 11–15. [In Russian]
- Shchegoleva NV, Ebel AL, Lashchinskiy NN, Ovchinnikov YuV (2019b) A new species of the genus *Ranunculus* from the Western Tien Shan. *Sistematicheskie Zametki po Materialam Gerbariya Imeni P.N. Krylova Tomskogo Gosudarstvennogo Universiteta* 119: 14–24. [In Russian] <https://doi.org/10.17223/20764103.119.2>
- Shennikov AP (1950) *Ecology of plants*. Soviet Science, Moscow, 375 pp. [In Russian]
- Smirnov SV, Kechaykin AA, Sinitsyna TA, Shmakov AI (2018) Synopsis of the genus *Saussurea* DC. (Asteraceae) of Eurasia. *Ukrainian Journal of Ecology* 8(4): 270–285.
- Soskov YD (2011) The genus *Calligonum* L. (systematics, geography, evolution, introduction). *RITS SibNSHB, Novosibirsk*, 362 pp. [In Russian]
- Stepanova EF (1962) Vegetation and flora of the Tarbagatai Ridge. *Academy of Sciences of Kazakhstan, Alma-Ata*, 433 pp. [In Russian]
- Suchorukow AP (2007) Zur Systematik und Chorologie der in Russland und den benachbarten Staaten (in den Grenzen der ehemaligen USSR) vorkommenden *Atriplex*-Arten (Chenopodiaceae). *Annalen des Naturhistorischen Museums in Wien, Serie B für Botanik und Zoologie* 108 B: 307–420.
- Sumbembayev AA, Lagus OA, Nowak S (2023) Seed morphometry of *Rheum* L. (Polygonaceae) species from Kazakhstan and its implications in taxonomy and species identification. *Biodiversitas (Surakarta)* 24(9): 4677–4692. <https://doi.org/10.13057/biodiv/d240908>
- Sitpayeva GT, Kudabayeva GM, Dimeyeva LA, Gemejiyeva NG, Vesselova PV (2020) Crop wild relatives of Kazakhstanian Tien Shan: Flora, vegetation, resources. *Plant Divers* 42: 19–32. <https://doi.org/10.1016/j.pld.2019.10.003>
- Takhtajan A (1969) Flowering plants. Origin and dispersal. Transl. by C. Jeffrey. Oliver & Boyd, Edinburgh, 310 pp. [In Russian]
- Thiers B (2023) Index Herbariorum: A Global Directory of Public Herbaria and Associated Staff. New York Botanical Garden's Virtual Herbarium. <http://sweetgum.nybg.org/science/ih/> [Accessed 25 March 2023]
- Tojibaev KS, Jang CG, Lazkov GA, Chang KS, Sitpayeva GT, Safarov N, Beshko NY, Muktubaeyeva SK, Vesselova PV, Turakulov I, Abdurakhmanova YY, Na NR, Park MS, Kyung C, Choi HJ, Oh BU, Oh SH (2020a) An annotated checklist of endemic vascular plants of the Tian-Shan Mountains in Central Asian countries. *Phytotaxa* 464(2): 117–158. <https://doi.org/10.11646/phytotaxa.464.2.1>

- Tojibaev K, Beshko N, Batoshov A, Azimova D, Yusupov Z, Deng T, Sun H (2020b) Flora of the Dzhizak Province, Uzbekistan. China Forestry Publishing House, 523 pp. [In Russian]
- Tojibaev K, Dekhkonov D, Ergashov I, Sun H, Deng T, Yusupov Z (2022) The synopsis of the genus *Tulipa* (Liliaceae) in Uzbekistan. *Phytotaxa* 573(2): 163–214. <https://doi.org/10.11646/phytotaxa.573.2.2>
- Tolmachev AI [Ed.] (1974) Endemic Highland Plants of the northern Asia. Science Publishers, Novosibirsk, 335 pp. [In Russian]
- Vaganov A (2023) Biodiversity of the Altai-Sayan ecoregion. Version 1.2. Altai State University. <https://doi.org/10.15468/6cvgov> [Accessed via GBIF.org on 2024-01-28]
- Vaganov A, Shmakov AI (2020) Biodiversity of Altai-Sayan Ecoregion Database of Plants and Fungi. Altai State University. <https://doi.org/10.15468/6cvgov> [Accessed via GBIF.org on 2020-08-21]
- Vesselova PV (2016) Biotopical distribution of small mammals in the steppe zone of the Zavolzhie. *Arid Ecosystems* 1(66): 31–37. [In Russian]
- Villaseñor JL, Ortiz E, Hernández-Flores MM (2023) The vascular plant species endemic or nearly endemic to Puebla, Mexico. *Botanical Sciences* 101(4): 1207–1221. <https://doi.org/10.17129/botsci.3299>
- Wu ZY, Raven PH, Hong DY [Eds] (2008) Flora of China, Vol. 11. Science Press, Beijing, and Missouri Botanical Garden Press, St. Louis, 622 pp.
- Yakovlev GP (2003) Leguminosae. In: Grubov VP (Ed.) Plants of Central Asia: Plant Collection from China and Mongolia, Vol. 8a. Science Publishers, Inc., Enfield, NH, 170 pp.
- Yegorina AV, Zinchenko YuK, Zinchenko ES (2003) Physical geography of the eastern Kazakhstan. Alfa-Press, Ust-Kamenogorsk, 184 pp. [In Russian]
- Yurtsev BA (1976) Life forms: one of the key subjects in botany. In: Serebryakova TI (Ed.) Problems of ecological morphology of plants. Science Publishers, Moscow, 9–44. [In Russian]
- Zhao Y, Chen YP, Yuan JC, Paton AJ, Nuraliev MS, Zhao F, Drew BT, Salmaki Y, Turginov OT, Sun M, Sennikov AN, Yu XQ, Li B, Xiang CL (2023) Museomics in Lamiaceae: Resolving the taxonomic mystery of *Pseudomarrubium*. *Current Plant Biology* 35–366: 100300. <https://doi.org/10.1016/j.cpb.2023.100300>
- Zolotukhin NI, Chkalov AV (2019) The genus *Alchemilla* L. (Rosaceae) in the Altai State Nature Reserve and adjacent areas. *Turczaninowia* 22(2): 5–42. [In Russian] <https://doi.org/10.14258/turczaninowia.22.2.1>
- Zonneveld BJ (2009) The systematic value of nuclear genome size for “all” species of *Tulipa* L. (Liliaceae). *Plant Systematics and Evolution* 281(1–4): 217–245. <https://doi.org/10.1007/s00606-009-0203-7>

Appendix 1. Annotated checklist of endemic taxa of Kazakhstan

In the checklist, families are listed in alphabetical order; lower-level taxa (genera, species and subspecies) within a family are also listed in alphabetical order.

The following information is given after the name of each taxon:

Life form (Lf) and life cycle (Lc) according to the Flora of Kazakhstan (Pavlov 1956); conservation status (Cs) according to the Red Book of Kazakhstan (Baitulin 2014). The species included in the Red Book of Kazakhstan are denoted by RB.

The distribution (D) of each taxon in Kazakhstan is given according to the floristic division of Kazakhstan (Pavlov 1956), where the territory of the country is divided into 29 floristic regions and seven subregions: 1 – Syrt, 2 – Tobol-Ishim,

3 – Irtysh, 4 – Semipalatinsk pine forest, 5 – Kokchetav, 6 – Caspian Region, 6a – Bukeev, 7 – Aktobe, 7a – Mugojary, 8 – Emba, 9 – Turgay, 10 – Western Upland, 10a – Ulutau, 11 – Eastern Upland, 11a – Karkaraly, 12 – Zaysan, 13 – Northern Ustyrt, 13a – Buzachi, 13b – Mangyshlak, 14 – Aral Region, 15 – Kyzylorda, 16 – Betpak-Dala, 17 – Moiynkum, 18 – Balkhash-Alakol, 19 – Southern Ustyrt, 20 – Kyzylkum, 21 – Turkestan, 22 – Altai, 23 – Tarbagatai, 24 – Hungarian Alatau, 25 – Trans-Ili Kungey Alatau, 25a – Ketmen-Terskey Alatau, 26 – Chu-Ili Range, 27 – Kyrgyz Alatau, 28 – Karatau, 29 – Western Tian Shan. Asterisks refer to annotations placed after the checklist.

Fam. 1. Amaranthaceae Juss.

Gen. 1. *Anabasis* L.

1. *Anabasis gypsicola* Iljin [Lf: Shrub. Lc: Per. D: 13, 16, 28]
2. *Anabasis turgaica* Iljin & Krasch. [Lf: Herb. Lc: Per. Cs: RB. D: 10a]

Gen. 2. *Arthrophytum* Schrenk

3. *Arthrophytum balchaschense* (Iljin) Botsch. [Lf: Subshrub. Lc: Per. D: 16, 17, 18, 25]
4. *Arthrophytum betpakdalense* Korovin & Mironov [Lf: Dwarf semishrub. Lc: Per. D: 16]
5. *Arthrophytum pulvinatum* Litv. [Lf: Dwarf semishrub. Lc: Per. D: 14]
6. *Arthrophytum subulifolium* Schrenk * [Lf: Subshrub. Lc: Per. D: 26]

Gen. 3. *Atriplex* L.

7. *Atriplex iljinii* Aellen * [Lf: Herb. Lc: An. D: 8, 13, 14]

Gen. 4. *Climacoptera* Botsch.

8. *Climacoptera turgaica* (Iljin) Botsch. [Lf: Herb. Lc: An. D: 2, 6, 9, 14]

Gen. 5. *Halimocnemis* C.A.Mey.

9. *Halimocnemis mironovii* Botsch. [Lf: Herb. Lc: An. D: 16, 26]

Gen. 6. *Horaninovia* Fisch. & C.A.Mey.

10. *Horaninovia capitata* Sukhor. [Lf: Herb. Lc: An. D: 18]

Gen. 7. *Nanophyton* Less.

11. *Nanophyton erinaceum* subsp. *karataviense* U.P.Pratov [Lf: Subshrub. Lc: Per. D: 28]

Gen. 8. *Petrosimonia* Bunge

12. *Petrosimonia hirsutissima* (Bunge) Iljin ex Pavlov [Lf: Herb. Lc: An. D: 9, 13, 15, 16, 17, 18]

Gen. 9. *Salsola* L.

13. *Salsola euryphylla* Botsch. [Lf: Undershrub. Lc: Per.Cs: RB. D: 14]

Gen. 10. *Suaeda* Forssk. ex J.F.Gmel.

14. *Suaeda scabra* Lomon. [Lf: Herb. Lc: An. D: 14]

Fam. 2. Amaryllidaceae J.St.-Hil.

Gen. 11. *Allium* L.

15. *Allium azutavicum* Kotukhov [Lf: Herb. Lc: Per. D: 22]
16. *Allium bajtulinii* Bajtenov & I.I.Kamenetskaya [Lf: Herb. Lc: Per. D: 25]
17. *Allium goloskokovii* Vved. [Lf: Herb. Lc: Per. D: 23, 24]
18. *Allium ivasczenkoae* Kotukhov [Lf: Herb. Lc: Per. D: 22]
19. *Allium iliense* Regel [Lf: Herb. Lc: Per. D: 26]
20. *Allium jaxarticum* Vved. [Lf: Herb. Lc: Per. D: 29]
21. *Allium kasteki* Popov. [Lf: Herb. Lc: Per. Cs: RB. D: 25]

22. *Allium kokuense* R.M. Fritsch, N. Friesen & S.V. Smirn. [Lf: Herb. Lc: Per. D: 24]
23. *Allium kujukense* Vved. [Lf: Herb. Lc: Per. D: 28, 29]
24. *Allium lasiophyllum* Vved. [Lf: Herb. Lc: Per. D: 25, 25a]
25. *Allium lehmannianum* Merckl. ex Bunge [Lf: Herb. Lc: Per. D: 9, 10, 15, 16, 20]
26. *Allium lepsicum* R.M.Fritsch, N.Friesen & S.V.Smirn. [Lf: Herb. Lc: Per. D: 24]
27. *Allium oreoprasoides* Vved. [Lf: Herb. Lc: Per. D: 28]
28. *Allium sergii* Vved. [Lf: Herb. Lc: Per. Cs: RB. D: 28]
29. *Allium scrobiculatum* Vved. [Lf: Herb. Lc: Per. D: 16, 17, 26]
30. *Allium subscabrum* (Regel) R.M.Fritsch [Lf: Herb. Lc: Per. D: 18, 26]
31. *Allium toksanbaicum* N.Friesen & Veselova [Lf: Herb. Lc: Per. D: 24]
32. *Allium turtschicum* Regel [Lf: Herb. Lc: Per. Cs: RB. D: 28]
33. *Allium victoris* Vved. [Lf: Herb. Lc: Per. D: 29]
34. *Allium viridulum* Ledeb. [Lf: Herb. Lc: Per. D: 11, 18, 23]
35. *Allium zaissanicum* Kotukhov [Lf: Herb. Lc: Per. D: 12]

Fam. 3. Apiaceae Lindl.

Gen. 12. *Autumnalia* Pimenov

36. *Autumnalia botschantzevii* Pimenov [Lf: Herb. Lc: Per. D: 28, 29]

Gen. 13. *Eryngium* L.

37. *Eryngium karatavicum* Iljin [Lf: Herb. Lc: Per. Cs: RB. D: 28]

Gen. 14. *Ferula* L.

38. *Ferula glaberrima* Korovin [Lf: Herb. Lc: Per. Cs: RB. D: 17]
39. *Ferula gypsacea* Korovin [Lf: Herb. Lc: Per. Cs: RB. D: 21]
40. *Ferula leucographa* Korovin * [Lf: Herb. Lc: Per. Cs: RB. D: 28]
41. *Ferula malacophylla* Pimenov & J.V.Baranova [Lf: Herb. Lc: Per. Cs: RB. D: 28]
42. *Ferula pachyphylla* Korovin [Lf: Herb. Lc: Per. D: 28]
43. *Ferula sugatensis* Bajtenov [Lf: Herb. Lc: Per. Cs: RB. D: 25]
44. *Ferula taucumica* Baitenov [Lf: Herb. Lc: Per. Cs: RB. D: 18]
45. *Ferula xeromorpha* Korovin [Lf: Herb. Lc: Per. Cs: RB. D: 21]

Gen. 15. *Hyalolaena* Bunge

46. *Hyalolaena tschuiensis* (Pavlov) Pimenov & Kljuykov [Lf: Herb. Lc: Per. Cs: RB. D: 26]

Gen. 16. *Karatavia* Pimenov & Lavrova

47. *Karatavia kultiassovii* (Korovin) Pimenov & Lavrova [Lf: Herb. Lc: Per. Cs: RB. D: 28, 29]

Gen. 17. *Pachypleurum* Ledeb.

48. *Pachypleurum altaicum* Revuschkin [Lf: Herb. Lc: Per. D: 22]

Gen. 18. *Pilopleura* Schischk.

49. *Pilopleura goloskokovii* (Korovin) Pimenov [Lf: Herb. Lc: Per. Cs: RB. D: 24]

Gen. 19. *Prangos* Lindl.

50. *Prangos dzhungarica* Pimenov [Lf: Herb. Lc: Per. D: 24]
51. *Prangos equisetoides* Kuzjmina [Lf: Herb. Lc: Per. Cs: RB. D: 28]
52. *Prangos lachnantha* (Korovin) Pimenov & Kljuykov [Lf: Herb. Lc: Per. Cs: RB. D: 17]
53. *Prangos multicostata* Kljuykov & Lyskov * [Lf: Herb. Lc: Per. D: 23]

Gen. 20. *Schrenkia* Fisch. & C.A.Mey.

54. *Schrenkia congesta* Korovin [Lf: Herb. Lc: Per. D: 28, 29]

55. *Schrenkia involucrata* Regel & Schmalh. [Lf: Herb. Lc: Per. D: 16, 17, 26, 28, 29]

56. *Schrenkia kultiassovii* Korovin [Lf: Herb. Lc: Per. Cs: RB. D: 29]

57. *Schrenkia papillaris* Regel & Schmalh. [Lf: Herb. Lc: Per. D: 28]

Gen. 21. *Schtschurowskia* Regel & Schmalh.

58. *Schtschurowskia margaritae* Korovin [Lf: Herb. Lc: Per. Cs: RB. D: 28]

Gen. 22. *Seseli* L.

59. *Seseli betpakdalense* Bajtenov [Lf: Herb. Lc: Per. D: 16]

60. *Seseli mironovii* (Korovin) Pimenov & Sdobnina [Lf: Herb. Lc: Per. D: 16]

Gen. 23. *Sphaenolobium* Pimenov

61. *Sphaenolobium korovinii* Pimenov & Kljuykov [Lf: Herb. Lc: Per. D: 28, 29]

Gen. 24. *Tschulaktavia* Bajtenov ex Pimenov & Kljuykov

62. *Tschulaktavia saxatilis* (Bajtenov) Bajtenov ex Pimenov & Kljuykov [Lf: Herb. Lc: Per. Cs: RB. D: 24]

Fam. 4. Asparagaceae Juss.

Gen. 25. *Asparagus* Tourn. ex L.

63. *Asparagus vvedenskyi* Botsch. [Lf: Herb. Lc: Per. Cs: RB. D: 21]

Fam. 5. Asteraceae Bercht. & J.Presl

Gen. 26. *Alfredia* Cass.

64. *Alfredia integrifolia* (Iljin) Tulyag. [Lf: Herb. Lc: Per. D: 24]

Gen. 27. *Amberboa* (Pers.) Less.

65. *Amberboa takhtajanii* Gabrieljan [Lf: Herb. Lc: An. D: 12]

Gen. 28. *Arctium* L.

66. *Arctium alberti* (Regel & Schmalh.) S.López, Romasch., Susanna & N.García [Lf: Herb. Lc: Per. D: 28, 29]

67. *Arctium arctiodes* (Schrenk) Kuntze [Lf: Herb. Lc: Per. D: 9, 10, 15, 16]

68. *Arctium grandifolium* (Kult.) S.López, Romasch., Susanna & N.García RB. [Lf: Herb. Lc: Per. Cs: RB. D: 28, 29]

69. *Arctium ugamense* (Karmysch.) S.López, Romasch., Susanna & N.García [Lf: Herb. Lc: Per. D: 29]

Gen. 29. *Artemisia* L.

70. *Artemisia aralensis* Krasch. [Lf: Dwarf semishrub. Lc: Per. D: 7, 9, 10, 14]

71. *Artemisia camelorum* Krasch. [Lf: Dwarf semishrub. Lc: Per. D: 7, 7a, 8, 9, 10, 14, 15]

72. *Artemisia filatovae* Kupr. [Lf: Dwarf semishrub. Lc: Per. D: 10]

73. *Artemisia hippolyti* A.Butkov [Lf: Dwarf semishrub. Lc: Per. D: 16]

74. *Artemisia kasakorum* (Krasch.) Pavlov [Lf: Herb. Lc: Per. D: 8, 11]

75. *Artemisia kotuchovii* Kupr. [Lf: Herb. Lc: Per. D: 22]

76. *Artemisia quinqueloba* Trautv. [Lf: Subshrub. Lc: Per. D: 7, 8, 14, 15, 16]

77. *Artemisia saurensis* Kupr. [Lf: Dwarf semishrub. Lc: Per. D: 23]

78. *Artemisia valida* Krasch. ex Poljakov [Lf: Dwarf semishrub. Lc: Per. D: 21, 28]

Gen. 30. *Brachanthemum* DC.

79. *Brachanthemum kasakorum* Krasch. [Lf: Dwarf semishrub. Lc: Per. D: 2, 10, 16, 18]

Gen. 31. *Cancriniella* Tzvelev

80. *Cancriniella krascheninnikovii* (Rubtzov) Tzvelev [Lf: Herb. Lc: Per. Cs: RB. D: 16, 26]

Gen. 32. *Centaurea* L.

81. *Centaurea kryloviana* Serg. * [Lf: Herb. Lc: Per. D: 11, 12, 22]

Gen. 33. *Chondrilla* L.

82. *Chondrilla bosseana* Iljin [Lf: Herb. Lc: Per. D: 17, 18]
 83. *Chondrilla macra* Iljin [Lf: Herb. Lc: Per. D: 16, 17]
 84. *Chondrilla mujunkumensis* Iljin & Igolkin [Lf: Herb. Lc: Per. D: 17, 18]

Gen. 34. *Cousinia* Cass.

85. *Cousinia aspera* (Kult.) Karmysch. [Lf: Herb. Lc: Per. D: 29]
 86. *Cousinia gomolitzkii* Juz. ex Tscherneva [Lf: Herb. Lc: Per. D: 28]
 87. *Cousinia kasachstanica* Sennikov [Lf: Herb. Lc: Per. D: 27]
 88. *Cousinia mindshelkensis* B.Fedtsch. [Lf: Herb. Lc: Per. Cs: RB. D: 28]
 89. *Cousinia perovskiensis* (Bornm.) Juz. ex Tschern. [Lf: Herb. Lc: Per. D: 14, 15, 16, 17, 18, 20, 24]
 90. *Cousinia schepsaica* Karmysch. [Lf: Herb. Lc: Per. D: 29]
 91. *Cousinia turkestanica* (Regel) Juz. [Lf: Herb. Lc: Per. D: 29]
 92. *Cousinia xanthiocephala* Tscherneva [Lf: Herb. Lc: Per. D: 21]

Gen. 35. *Echinops* L.

93. *Echinops kasakorum* Pavlov [Lf: Herb. Lc: Per. Cs: RB. D: 28]
 94. *Echinops pubisquameus* Iljin [Lf: Herb. Lc: Per. D: 28]
 95. *Echinops subglaber* Schrenk [Lf: Herb. Lc: Per. D: 15, 16, 28]
 96. *Echinops transiliensis* Golosk. [Lf: Herb. Lc: Per. D: 25, 26]

Gen. 36. *Galatella* Cass.

97. *Galatella bectauatensis* Kupr. & Koroljuk [Lf: Herb. Lc: Per. D: 11]
 98. *Galatella polygaloides* Novopokr. [Lf: Herb. Lc: Per. D: 25a]
 99. *Galatella saxatilis* Novopokr. [Lf: Herb. Lc: Per. Cs: RB. D: 25]

Gen. 37. *Hieracium* L.

100. *Hieracium bectauatense* Kupr. [Lf: Herb. Lc: Per. D: 11]

Gen. 38. *Jurinea* Cass.

101. *Jurinea almaatensis* Iljin [Lf: Herb. Lc: Per. Cs: RB. D: 25]
 102. *Jurinea bracteata* Regel & Schmalh. [Lf: Herb. Lc: Per. D: 24]
 103. *Jurinea cephalopoda* Iljin [Lf: Herb. Lc: Per. Cs: RB. D: 15, 21, 28]
 104. *Jurinea czilikinoana* Iljin [Lf: Herb. Lc: Per. D: 28]
 105. *Jurinea eximia* Tekutj. [Lf: Herb. Lc: Per. Cs: RB. D: 28]
 106. *Jurinea fedtschenkoana* Iljin [Lf: Herb. Lc: Per. Cs: RB. D: 1, 7, 7a, 10]
 107. *Jurinea hamulosa* Rubtzov [Lf: Herb. Lc: Per. D: 23, 24]
 108. *Jurinea karatavica* Iljin [Lf: Herb. Lc: Per. D: 17, 18]
 109. *Jurinea kazachstanica* Iljin [Lf: Herb. Lc: Per. D: 6, 7, 8, 14]
 110. *Jurinea knorringiana* Iljin [Lf: Herb. Lc: Per. D: 28]
 111. *Jurinea krascheninnikovii* Iljin [Lf: Herb. Lc: Per. D: 9, 11a, 15, 16, 17, 26]
 112. *Jurinea lithophila* Rubtzov [Lf: Herb. Lc: Per. D: 24]
 113. *Jurinea monticola* Iljin [Lf: Herb. Lc: Per. D: 28]
 114. *Jurinea multiceps* Iljin [Lf: Herb. Lc: Per. D: 28]
 115. *Jurinea pineticola* Iljin [Lf: Herb. Lc: Per. D: 3, 4]
 116. *Jurinea rhizomatoidea* Iljin [Lf: Herb. Lc: Per. D: 28]
 117. *Jurinea robusta* Schrenk [Lf: Herb. Lc: Per. Cs: RB. D: 16, 25, 26]
 118. *Jurinea serratuloides* Iljin [Lf: Herb. Lc: Per. D: 12, 23]
 119. *Jurinea sudunensis* Korsh. [Lf: Herb. Lc: Per. D: 12, 18, 24]
 120. *Jurinea xerophytica* Iljin [Lf: Herb. Lc: Per. D: 4, 9, 11, 12, 14]

Gen. 39. *Lamyropsis* (Kharadze) Dittrich

121. *Lamyropsis macracantha* (Schrenk) Dittrich [Lf: Herb. Lc: Per. D:23, 24]
- Gen. 40. *Lepidolopha* C.Winkl.**
122. *Lepidolopha gomolitzkii* Kovalevsk. & Safral. [Lf: Herb. Lc: Per. D: 28]
123. *Lepidolopha karatavica* Pavlov [Lf: Herb. Lc: Per. Cs: RB. D: 28]
124. *Lepidolopha krascheninnikovii* Czil. ex Kovalevsk. & Safral. [Lf: Herb. Lc: Per. D: 28]
125. *Lepidolopha talassica* Kovalevsk. & Safral. [Lf: Herb. Lc: Per. RB. D: 29]
- Gen. 41. *Rhaponticum* Vaill.**
126. *Rhaponticum karatavicum* Regel & Schmalh. [Lf: Herb. Lc: Per. Cs: RB. D: 28]
- Gen. 42. *Ligularia* Cass.**
127. *Ligularia pavlovii* (Lipsch.) Cretz. [Lf: Herb. Lc: Per. Cs: RB. D: 28]
- Gen. 43. *Pseudoglossanthis* Poljakov**
128. *Pseudoglossanthis arctodshungarica* (Golosc.) Kamelin [Lf: Subshrub. Lc: Per. Cs: RB. D: 24]
129. *Pseudoglossanthis simulans* (Pavlov) Kamelin [Lf: Herb. Lc: Per. D: 29]
- Gen. 44. *Pseudopodospermum* (Lipsch. &Krasch.) Kuth.**
130. *Pseudopodospermum chantavicum* (Pavlov) Zaika, Sukhor. & N.Kilian [Lf: Herb. Lc: Per. Cs: RB. D: 26]
- Gen. 45. *Rhaponticoides* Vaill.**
131. *Rhaponticoides kultiassovii* (Iljin) Negaresh [Lf: Herb. Lc: Per. Cs: RB. D: 28]
132. *Rhaponticoides phyllopoda* (Iljin) Negaresh [Lf: Herb. Lc: Per. D: 28, 29]
133. *Rhaponticoides zaissanica* Kupr., A.L. Ebel et Khrustaleva [Lf: Herb. Lc: Per. D: 12]
- Gen. 46. *Saussurea* DC.**
134. *Saussurea mikeschinii* Iljin [Lf: Subshrub. Lc: Per. Cs: RB. D: 28]
135. *Saussurea ninae* Iljin [Lf: Herb. Lc: Per. D: 24]
136. *Saussurea pseudoblada* Lipsch. ex Filat. [Lf: Herb. Lc: Per. D: 24]
- Gen. 47. *Scorzonera* L.**
137. *Scorzonera dianthoides* (Lipsch. & Krasch.) Lipsch. [Lf: Herb. Lc: Per. D: 11]
138. *Scorzonera franchetii* Lipsch. [Lf: Herb. Lc: Per. D: 29]
139. *Scorzonera vavilovii* Kult. [Lf: Dwarf semishrub. Lc: Per. D: 28, 29]
- Gen. 48. *Senecio* L.**
140. *Senecio iljinii* Schischk. [Lf: Herb. Lc: Per. D: 24]
141. *Senecio nuraniae* Roldugin [Lf: Herb. Lc: An. D: 29]
- Gen. 49. *Takhtajiantha* Nazarova**
142. *Takhtajiantha veresczaginii* (Kamelin & S.V.Smirn.) Zaika, Sukhor. & N.Kilian [Lf: Herb. Lc: Per. D: 22]
- Gen. 50. *Tanacetopsis* (Tzvelev) Kovalevsk.**
143. *Tanacetopsis goloskokovii* (Poljakov) Karmysch. [Lf: Herb. Lc: Per. Cs: RB. D: 24, 25]
144. *Tanacetopsis pjataevae* (Kovalevsk.) Karmysch. [Lf: Herb. Lc: Per. Cs: RB. D: 28]
145. *Tanacetopsis popovii* Kamelin & Kovalevsk. [Lf: Herb. Lc: Per. D: 28]
- Gen. 51. *Tanacetum* L.**
146. *Tanacetum corymbiforme* (Tzvelev) K.Bremer & Humphries [Lf: Herb. Lc: Per. D: 23, 24]
147. *Tanacetum kelleri* (Krylov & Plotn.) Takht. [Lf: Herb. Lc: Per. Cs: RB. D: 22]
148. *Tanacetum mindshelkense* Kovalevsk. [Lf: Herb. Lc: Per. D: 28]

149. *Tanacetum saryarkense* Kamelin [Lf: Herb. Lc: Per. D: 16, 26]
 150. *Tanacetum saxicola* (Krasch.) Tzvelev [Lf: Herb. Lc: Per. Cs: RB. D: 7a, 8]
 151. *Tanacetum ulutavicum* Tzvelev [Lf: Herb. Lc: Per. Cs: RB. D: 10a]

Gen. 52. *Taraxacum* F.H.Wigg.

152. *Taraxacum almaatense* Schischk. [Lf: Herb. Lc: Per. D: 25]
 153. *Taraxacum arasanum* R.Doll [Lf: Herb. Lc: Per. D: 25]
 154. *Taraxacum atrochlorinum* Kirschner & Štěpánek [Lf: Herb. Lc: Per. D: 25]
 155. *Taraxacum botschantzevii* Schischk. [Lf: Herb. Lc: Per. D: 28]
 156. *Taraxacum cornucopiae* Kirschner & Štěpánek [Lf: Herb. Lc: Per. D: 25]
 157. *Taraxacum corvinum* Kirschner & Štěpánek [Lf: Herb. Lc: Per. D: 25]
 158. *Taraxacum dzhungaricola* Kirschner & Štěpánek [Lf: Herb. Lc: Per. D: 24]
 159. *Taraxacum glabellum* Schischk. [Lf: Herb. Lc: Per. D: 28]
 160. *Taraxacum karatavicum* Pavlov [Lf: Herb. Lc: Per. D: 28]
 161. *Taraxacum kasachiforme* R.Doll [Lf: Herb. Lc: Per. D: 25]
 162. *Taraxacum kasachum* R.Doll [Lf: Herb. Lc: Per. D: 25]
 163. *Taraxacum magnum* Korol. [Lf: Herb. Lc: Per. D: 25a]
 164. *Taraxacum medeense* R.Doll [Lf: Herb. Lc: Per. D: 24, 25]
 165. *Taraxacum perpusillum* Schischk. [Lf: Herb. Lc: Per. D: 24]
 166. *Taraxacum pseudolugubre* R.Doll [Lf: Herb. Lc: Per. D: 25]
 167. *Taraxacum pseudotianschanicum* R.Doll [Lf: Herb. Lc: Per. D: 8]
 168. *Taraxacum saposhnikovii* Schischk. [Lf: Herb. Lc: Per. D: 23, 24]
 169. *Taraxacum sublilacinum* Kirschner & Štěpánek [Lf: Herb. Lc: Per. D: 25]
 170. *Taraxacum urdzhareense* Orazova [Lf: Herb. Lc: Per. D: 23]
 171. *Taraxacum violaceum* R.Doll [Lf: Herb. Lc: Per. D: 25]

Gen. 53. *Tragopogon* L.

172. *Tragopogon karelinii* S.A.Nikitin [Lf: Herb. Lc: Bi. D: 10, 16, 18, 23, 24]

Gen. 54. *Vickifunkia* C.Ren, L.Wang, I.D.Illar. & Q.E.Yang

173. *Vickifunkia kareliniana* (Stschegl.) C.Ren, L.Wang, I.D.Illar. & Q.E.Yang
 [Lf: Herb. Lc: Per. D: 23]

Fam. 6. Berberidaceae Juss.

Gen. 55. *Berberis* L.

174. *Berberis karkaralensis* Kornil. & Potapov [Lf: Shrub. Lc: Per. Cs: RB. D: 11a]

Fam. 7. Betulaceae Gray

Gen. 56. *Betula* L.

175. *Betula karagandensis* V.N.Vassil. [Lf: Tree. Lc: Per. D: 11a]
 176. *Betula saviczii* V.N.Vassil. [Lf: Tree. Lc: Per. D: 10]

Fam. 8. Bignoniaceae Juss.

Gen. 57. *Incarvillea* Juss.

177. *Incarvillea semiretschenskia* (B.Fedtsch.) Grierson [Lf: Herb. Lc: Per. Cs: RB. D: 26]

Fam. 9. Boraginaceae Juss.

Gen. 58. *Eritrichium* Schrad. ex Gaudin

178. *Eritrichium relictum* Kudab. [Lf: Herb. Lc: Per. D: 24]

Gen. 59. *Heliotropium* Tourn. ex L.

179. *Heliotropium parvulum* Popov [Lf: Herb. Lc: An. Cs: RB. D: 16, 18, 25]

Gen. 60. *Lappula* Moench

- 180. *Lappula baitenovii* Kudab. [Lf: Herb. Lc: Bi. D: 25]
- 181. *Lappula coronifera* Popov [Lf: Herb. Lc: An. D: 11]
- 182. *Lappula cristata* (Bunge) B.Fedtsch. *[Lf: Herb. Lc: An. D: 10, 11, 12]
- 183. *Lappula diploloma* (Fisch. & C.A.Mey.) Gürke [Lf: Herb. Lc: An. D: 9, 11]
- 184. *Lappula glabrata* Popov [Lf: Herb. Lc: Bi. Cs: RB. D: 16]
- 185. *Lappula ketmenica* Kudab. [Lf: Herb. Lc: An. D: 25a]
- 186. *Lappula kuprijanovii* Ovczinnikova [Lf: Herb. Lc: Bi. RB. D: 28]
- 187. *Lappula lipschitzii* Popov [Lf: Herb. Lc: An. D: 21]
- 188. *Lappula pavlovii* Golosk. [Lf: Herb. Lc: An. D: 24]
- 189. *Lappula saphronovae* Kamelin [Lf: Herb. Lc: Bi. D: 13b]
- 190. *Lappula zaissanica* (Aralbaev) Aralbaev [Lf: Herb. Lc: An. D: 12]

Gen. 61. *Lepechiniella* Popov

- 191. *Lepechiniella austrodshungarica* Golosk. [Lf: Herb. Lc: An.-Bi. D: 10, 24]
- 192. *Lepechiniella michaelis* Golosk. [Lf: Herb. Lc: Per. Cs: RB. D: 24]
- 193. *Lepechiniella omphaloides* (Schrenk) Popov [Lf: Herb. Lc: Bi. D: 10]
- 194. *Lepechiniella saurica* (Bajtenov & Kudab.) Ovczinnikova [Lf: Herb. Lc: An. D: 23]

Gen. 62. *Mattiastrum* (Boiss.) Brand

- 195. *Mattiastrum karataviense* (Pavlov ex Popov) Czerep. [Lf: Herb. Lc: Per. Cs: RB. D: 28]

Gen. 63. *Myosotis* L.

- 196. *Myosotis kazakhstanica* O.D.Nikif. [Lf: Herb. Lc: An. D: 10, 10a, 11, 11a]

Gen. 64. *Paracaryum* Boiss.

- 197. *Paracaryum integerrimum* Myrz. [Lf: Herb. Lc: Per. Cs: RB. D: 28]

Gen. 65. *Rindera* Pall.

- 198. *Rindera ochroleuca* Kar. & Kir. * [Lf: Herb. Lc: Per. Cs: RB. D: 18]

Gen. 66. *Rochelia* Rchb.

- 199. *Rochelia leiosperma* (Popov) Golosk. [Lf: Herb. Lc: An. D: 24]

Gen. 67. *Sauria* Bajtenov

- 200. *Sauria akkolia* Bajtenov [Lf: Herb. Lc: Per. D: 17]

Fam. 10. Brassicaceae Burnett

Gen. 68. *Botschantzevia* Nabiev

- 201. *Botschantzevia karatavica* (Lipsch. & Pavlov) Nabiev [Lf: Dwarf semishrub. Lc: Per. Cs: RB. D: 28]

Gen. 69. *Clausia* Korn-Trotzky.

- 202. *Clausia kasakorum* Pavlov [Lf: Herb. Lc: Per. D: 10a]
- 203. *Clausia robusta* Pachom. [Lf: Herb. Lc: Per. D: 5]

Gen. 70. *Erysimum* Tourn. ex L.

- 204. *Erysimum kazachstanicum* Botsch. [Lf: Herb. Lc: Bi. D: 10, 10a, 11, 23]

Gen. 71. *Eutrema* R.Br.

- 205. *Eutrema halophilum* (C.A.Mey.) Al-Shehbaz & Warwick* [Lf: Herb. Lc: An. D: 2, 3, 4, 11, 12]
- 206. *Eutrema platypetalum* (Schrenk) Al-Shehbaz & Warwick [Lf: Herb. Lc: Per. D: 24]

Gen. 72. *Isatis* Tourn. ex L.

- 207. *Isatis canaliculata* (Vassilcz.) V.V.Botschantz. [Lf: Herb. Lc: Bi. D: 9, 10]
- 208. *Isatis deserti* (N.Busch) V.V.Botschantz. [Lf: Herb. Lc: An. D: 16]

Gen. 73. *Lepidium* L.

- 209. *Lepidium jarmolenkoi* V.M.Vinogr. [Lf: Herb. Lc: Per. D: 16]
- 210. *Lepidium karataviense* Regel & Schmalh. [Lf: Herb. Lc: Per. D: 28]
- 211. *Lepidium mummenhoffianum* Al-Shehbaz [Lf: Herb. Lc: Per. D: 24]
- 212. *Lepidium pavlovii* Al-Shehbaz & Mummenhoff [Lf: Herb. Lc: Per. Cs: RB. D: 28, 29]
- 213. *Lepidium robustum* (Pavlov) Al-Shehbaz [Lf: Herb. Lc: Per. Cs: RB. D: 28]
- 214. *Lepidium sagittatum* (Kar. & Kir.) Al-Shehbaz [Lf: Herb. Lc: Per. Cs: RB. D: 23, 24]
- 215. *Lepidium trautvetteri* (Botsch.) Al-Shehbaz [Lf: Herb. Lc: Per. Cs: RB. D: 16, 18]

Gen. 74. *Parrya* R.Br.

- 216. *Parrya longicarpa* Krasn. [Lf: Herb. Lc: Per. D: 26]
- 217. *Parrya papillosa* (Vassilcz.) D.A.German & Al-Shehbaz [Lf: Herb. Lc: An. D: 28, 29]
- 218. *Parrya pavlovii* A.N.Vassiljeva [Lf: Herb. Lc: Per. D: 28]
- 219. *Parrya pazijae* (Pachom.) D.A.German & Al-Shehbaz [Lf: Undershrub. Lc: Per. D: 28, 29]
- 220. *Parrya saurica* (Pachom.) D.A.German & Al-Shehbaz [Lf: Herb. Lc: Per. D: 23]
- 221. *Parrya vvedenskyi* (Pachom.) D.A.German & Al-Shehbaz [Lf: Herb. Lc: Bi. D: 28, 29]

Gen. 75. *Rhammatophyllum* O.E.Schulz

- 222. *Rhammatophyllum pachyrhizum* (Kar. & Kir.) O.E.Schulz [Lf: Dwarf semishrub. Lc: Per. D: 7a, 8, 9, 10, 10a, 11, 13, 14, 16, 22, 24]

Gen. 76. *Strigosella* Boiss.

- 223. *Strigosella myrzakulovii* Bajtenov [Lf: Herb. Lc: An. D: 28, 29]

Fam. 11. Campanulaceae Juss.

Gen. 77. *Sergia* Fed.

- 224. *Sergia sewerzowii* (Regel) Fed. [Lf: Herb. Lc: Per. D: 28, 29]

Fam. 12. Caryophyllaceae Juss.

Gen. 78. *Eremogone* Fenzl

- 225. *Eremogone turlanica* (Bajtenov) Czerep. [Lf: Herb. Lc: Per. Cs: RB. D: 28]

Gen. 79. *Dianthus* L.

- 226. *Dianthus karataviensis* Pavlov [Lf: Herb. Lc: Per. RB. D: 28, 29]
- 227. *Dianthus multisquameus* Bondarenko & R.M.Vinogr. [Lf: Herb. Lc: Per. D: 28, 29]

Gen. 80. *Gypsophila* L.

- 228. *Gypsophila aulieatensis* B.Fedtsch. [Lf: Herb. Lc: Per. Cs: RB. D: 28]

Gen. 81. *Silene* L.

- 229. *Silene anisoloba* Schrenk [Lf: Herb. Lc: Per. D: 10, 10a]
- 230. *Silene betpakdalensis* Bajtenov [Lf: Herb. Lc: Per. Cs: RB. D: 13, 16, 26]
- 231. *Silene jaxartica* Pavlov [Lf: Herb. Lc: Per. Cs: RB. D: 28]
- 232. *Silene muslimii* Pavlov [Lf: Herb. Lc: Per. Cs: RB. D: 24, 26]

Fam. 13. Convolvulaceae Juss.

Gen. 82. *Cuscuta* L.

- 233. *Cuscuta camelorum* Pavlov [Lf: Herb. Lc: An. D: 28]

234. *Cuscuta elpassiana* Pavlov [Lf: Herb. Lc: An. D: 26]

235. *Cuscuta karatavica* Pavlov [Lf: Herb. Lc: An. D: 28]

Fam. 14. Crassulaceae J.St.-Hil.

Gen. 83. *Pseudosedum* (Boiss.) A.Berger

236. *Pseudosedum karatavicum* Boriss. [Lf: Herb. Lc: Per. Cs: RB. D: 28]

Fam. 15. Cyperaceae Juss.

Gen. 84. *Cyperus* L.

237. *Cyperus soongoricus* Kar. & Kir. [Lf: Herb. Lc: An. D: 12]

Fam. 16. Euphorbiaceae Juss.

Gen. 85. *Euphorbia* L.

238. *Euphorbia heptapotamica* Golosk. [Lf: Herb. Lc: An. D: 24]

239. *Euphorbia kalbaensis* Baikov & I.V.Khan [Lf: Herb. Lc: Per. D: 22]

240. *Euphorbia saurica* Baikov [Lf: Herb. Lc: Per. D: 23]

241. *Euphorbia yaroslavii* Poljakov [Lf: Herb. Lc: Per. Cs: RB. D: 25]

Fam. 17. Fabaceae Lindl.

Gen. 86. *Astragalus* L.

242. *Astragalus abbreviatus* Kar. & Kir. [Lf: Herb. Lc: Per. D: 24, 25, 25a, 26, 27]

243. *Astragalus arganaticus* Bunge [Lf: Herb. Lc: Per. D: 18, 24]

244. *Astragalus balchaschensis* Sumnev. [Lf: Herb. Lc: Per. D: 18]

245. *Astragalus brotherusii* Podlech [Lf: Herb. Lc: Per. D: 25]

246. *Astragalus chaetolobus* Bunge [Lf: Subshrub. Lc: Per. D: 4, 11, 22]

247. *Astragalus citoinflatus* Bondarenko [Lf: Herb. Lc: Per. D: 17]

248. *Astragalus clausii* C.A.Mey. [Lf: Herb. Lc: Per. D: 6]

249. *Astragalus cytisoides* Bunge [Lf: Dwarf semishrub. Lc: Per. D: 18]

250. *Astragalus fragiformis* Willd. [Lf: Undershrub. Lc: Per. D: 22]

251. *Astragalus georgii* Gontsch. [Lf: Dwarf semishrub. Lc: Per. D: 28]

252. *Astragalus inflatus* DC. [Lf: Undershrub. Lc: Per. D: 22]

253. *Astragalus jaxarticus* Pavlov [Lf: Herb. Lc: Per. D: 26, 28]

254. *Astragalus juvenalis* Delile [Lf: Herb. Lc: An. D: 10, 18]

255. *Astragalus karataviensis* Pavlov [Lf: Dwarf semishrub. Lc: Per. Cs: RB. D: 21, 28]

256. *Astragalus karatjubeki* Golosk. [Lf: Subshrub. Lc: Per. D: 16, 18]

257. *Astragalus kasachstanicus* subsp. *coloratus* Knjaz. [Lf: Herb. Lc: Per. D: 10, 11a]

258. *Astragalus kazymbeticus* Saposhn. ex Sumnev. [Lf: Herb. Lc: Per. D: 24]

259. *Astragalus kopalensis* Lipsky [Lf: Shrub. Lc: Per. Cs: RB. D: 24]

260. *Astragalus krascheninnikovii* Kamelin [Lf: Shrub. Lc: Per. Cs: RB. D: 16]

261. *Astragalus krasnovii* Popov [Lf: Herb. Lc: Per. D: 26]

262. *Astragalus leucocalyx* Popov * [Lf: Shrub. Lc: Per. D: 28]

263. *Astragalus lipschitzii* Pavlov [Lf: Undershrub. Lc: Per. D: 28, 29]

264. *Astragalus mokeevae* Popov [Lf: Subshrub. Lc: Per. D: 28]

265. *Astragalus neopopovii* Golosk. [Lf: Herb. Lc: Per. D: 24]

266. *Astragalus psammophilus* Golosk. [Lf: Subshrub. Lc: Per. D: 18]

267. *Astragalus pseudocytisoides* Popov Lf: Dwarf semishrub. Lc: Per. Cs: RB. D: 25, 26]

268. *Astragalus psilopus* Schrenk [Lf: Herb. Lc: Per. D: 18, 24]
269. *Astragalus pulposus* Popov [Lf: Herb. Lc: Per. D: 25]
270. *Astragalus pycnolobus* Bunge [Lf: Subshrub. Lc: Per. D: 12, 22]
271. *Astragalus rariflorus* Ledeb. * [Lf: Herb. Lc: Per. D: 11]
272. *Astragalus rubtzovii* Boriss. [Lf: Herb. Lc: Per. Cs: RB. D: 25a]
273. *Astragalus saphronovae* Kulikov [Lf: Dwarf semishrub. Lc: Per. D: 7, 13, 13b]
274. *Astragalus sarchanensis* Gontsch. [Lf: Herb. Lc: Per. D: 24]
275. *Astragalus semenovii* Bunge [Lf: Herb. Lc: Per. D: 18, 24, 25a]
276. *Astragalus sisyrodytes* Bunge [Lf: Herb. Lc: Per. D: 28]
277. *Astragalus spartioides* Kar. & Kir. [Lf: Subshrub. Lc: Per. D: 18]
278. *Astragalus speciosissimus* Pavlov [Lf: Subshrub. Lc: Per. D: 28]
279. *Astragalus subcaracugensis* Sitpaeva [Lf: Subshrub. Lc: Per. D: 9]
280. *Astragalus subternatus* Pavlov [Lf: Herb. Lc: Per. Cs: RB. D: 28]
281. *Astragalus sumnevicii* Pavlov [Lf: Herb. Lc: Per. Cs: RB. D: 16]
282. *Astragalus terekensis* Fisjun [Lf: Herb. Lc: Per. D: 24]
283. *Astragalus transnominatus* M.N.Abdull. [Lf: Herb. Lc: Per. D: 26, 28]
284. *Astragalus tscharynensis* Popov [Lf: Undershrub. Lc: Per. Cs: RB. D: 24, 25]
285. *Astragalus turajgyricus* Golosk. [Lf: Herb. Lc: Per. D: 25]
286. *Astragalus unilateralis* Kar. & Kir. [Lf: Herb. Lc: Per. D: 4, 7, 8, 11, 12, 22]
287. *Astragalus virens* Pavlov [Lf: Herb. Lc: Per. D: 28]

Gen. 87. Caragana Lam.

288. *Caragana media* Sanchir [Lf: Shrub. Lc: Per. D: 10, 11]

Gen. 88. Chesneya Lindl. ex Endl.

289. *Chesneya karatavica* Kamelin [Lf: Herb. Lc: Per. D: 28]

Gen. 89. Hedysarum L.

290. *Hedysarum bectauatavicum* Bajtenov [Lf: Herb. Lc: Per. Cs: RB. D: 11]
291. *Hedysarum chantavicum* Popov ex Bajtenov [Lf: Herb. Lc: Per. D: 26]
292. *Hedysarum karataviense* B.Fedtsch. [Lf: Herb. Lc: Per. Cs: RB. D: 28]
293. *Hedysarum nikolai* Kovalevsk. [Lf: Herb. Lc: Per. D: 28]
294. *Hedysarum mindshilkense* Bajtenov [Lf: Herb. Lc: Per. Cs: RB. D: 28]
295. *Hedysarum pallidiflorum* Pavlov [Lf: Herb. Lc: Per. D: 28]
296. *Hedysarum pavlovii* Bajtenov [Lf: Herb. Lc: Per. D: 28]
297. *Hedysarum tarbagataicum* Knjaz. [Lf: Herb. Lc: Per. D: 3, 11, 12, 22, 23]
298. *Hedysarum ulutavicum* Knjaz. [Lf: Herb. Lc: Per. D: 10a]
299. *Hedysarum villosissimum* Knjaz. [Lf: Herb. Lc: Per. D: 10, 11]

Gen. 90. Onobrychis Mill.

300. *Onobrychis alatavica* Bajtenov * [Lf: Herb. Lc: Per. Cs: RB. D: 25]

Gen. 91. Oxytropis DC.

301. *Oxytropis alberti-regelii* Vassilcz. [Lf: Herb. Lc: Per. D: 29]
302. *Oxytropis almaatensis* Bajtenov [Lf: Herb. Lc: Per. Cs: RB. D: 25, 25a]
303. *Oxytropis bajtulinii* Kotukhov [Lf: Herb. Lc: Per. D: 22]
304. *Oxytropis biloba* Saposhn. [Lf: Herb. Lc: Per. Cs: RB. D: 23]
305. *Oxytropis bosculensis* Golosk. [Lf: Herb. Lc: Per. D: 25]
306. *Oxytropis brevicaulis* Ledeb. [Lf: Herb. Lc: Per. D: 2, 3, 9, 10, 11, 18]
307. *Oxytropis canopatula* Vassilcz. [Lf: Herb. Lc: Per. D: 28]
308. *Oxytropis cretacea* Basil. [Lf: Herb. Lc: Per. D: 1]
309. *Oxytropis echidna* Vved. [Lf: Undershrub. Lc: Per. Cs: RB. D: 28]
310. *Oxytropis fruticulosa* Bunge [Lf: Undershrub. Lc: Per. D: 24]

311. *Oxytropis gebleriana* Schrenk [Lf: Herb. Lc: Per. D: 1, 2, 3, 7a, 9, 10, 11, 16, 18]
312. *Oxytropis heteropoda* Bunge [Lf: Herb. Lc: Per. D: 24, 25]
313. *Oxytropis karataviensis* Pavlov [Lf: Herb. Lc: Per. Cs: RB. D: 28]
314. *Oxytropis kyziltalensis* Vassilcz. [Lf: Herb. Lc: Per. D: 24]
315. *Oxytropis niedzweckiana* Popov [Lf: Herb. Lc: Per. Cs: RB. D: 25]
316. *Oxytropis pulvinoides* Vassilcz. [Lf: Herb. Lc: Per. D: 24]
317. *Oxytropis satpaevii* Bajtenov [Lf: Herb. Lc: Per. D: 11]
318. *Oxytropis subcapitata* Gontsch [Lf: Herb. Lc: Per. D: 28]
319. *Oxytropis subverticillaris* C.A.Mey. [Lf: Herb. Lc: Per. Cs: RB. D: 3, 10, 10a, 11]
320. *Oxytropis sumneviczii* Krylov [Lf: Herb. Lc: Per. D: 22]
321. *Oxytropis talgarica* Popov [Lf: Herb. Lc: Per. D: 25]
322. *Oxytropis tomentosa* Gontsch. [Lf: Herb. Lc: Per. D: 28]

Fam. 18. Frankeniaceae Desv.

Gen. 92. Frankenia L.

323. *Frankenia bucharica* subsp. *mironovii* (Botsch.) Chrtek [Lf: Subshrub. Lc: Per. D: 16, 25, 26]
324. *Frankenia bucharica* subsp. *transkaratavica* (Botsch.) Chrtek [Lf: Subshrub. Lc: Per. D: 17]

Fam. 19. Gentianaceae Juss.

Gen. 93. Comastoma Toyok.

325. *Comastoma irinae* (Pachom.) Czerep. [Lf: Herb. Lc: An. D: 25]

Fam. 20. Lamiaceae Martinov

Gen. 94. Dracocephalum L.

326. *Dracocephalum pavlovii* Roldugin [Lf: Dwarf semishrub. Lc: Per. D: 29]

Gen. 95. Phlomoides Moench.

327. *Phlomoides affinis* (Schrenk) Salmaki [Lf: Herb. Lc: Per. D: 10, 10a, 11, 16, 17, 18, 28]
328. *Phlomoides boraldaica* A.L.Ebel [Lf: Herb. Lc: Per. D: 28, 29]
329. *Phlomoides czuiliensis* (Golosc.) Adylov, Kamelin & Makhm. [Lf: Herb. Lc: Per. D: 26]
330. *Phlomoides eremostachydioides* (Popov) Y.Zhao & C.L.Xiang [Lf: Herb. Lc: Cs: RB. Per. D: 28]
331. *Phlomoides gymnocalyx* (Schrenk) Adylov, Kamelin & Makhm. [Lf: Herb. Lc: Per. D: 18, 24, 26]
332. *Phlomoides iliensis* (Regel) Adylov, Kamelin & Makhm [Lf: Herb. Lc: Per. D: 18, 24]
333. *Phlomoides pectinata* (Popov) Adylov, Kamelin & Makhm. [Lf: Herb. Lc: Per. D: 28]
334. *Phlomoides rotala* (Schrenk ex Fisch., C.A.Mey. & Avé-Lall.) Salmaki [Lf: Herb. Lc: Per. D: 18]
335. *Phlomoides septentrionalis* (Popov) Adylov, Kamelin & Makhm. [Lf: Herb. Lc: Per. D: 27, 28, 29]

Gen. 96. Phlomis L.

336. *Phlomis mindshelkensis* Lazkov [Lf: Herb. Lc: Per. D: 28]

Gen. 97. *Lagochilus* Bunge ex Benth.

337. *Lagochilus androssowii* Knorring [Lf: Dwarf semishrub. Lc: Per. D: 15, 28]
 338. *Lagochilus longidentatus* Knorring [Lf: Dwarf semishrub. Lc: Per. D: 16, 28]
 339. *Lagochilus taukumensis* Tzukerv. [Lf: Dwarf semishrub. Lc: Per. D: 18]

Gen. 98. *Leonurus* L.

340. *Leonurus incanus* V.I.Krecz. & Kuprian. [Lf: Herb. Lc: Per. D: 24]

Gen. 99. *Salvia* L.

341. *Salvia trautvetteri* Regel [Lf: Herb. Lc: Per. D: 28, 29]

Gen. 100. *Scutellaria* L.

342. *Scutellaria androssovii* Juz. [Lf: Herb. Lc: Per. D: 15, 28]
 343. *Scutellaria karatavica* Juz. [Lf: Herb. Lc: Per. Cs: RB. D: 28]
 344. *Scutellaria kurssanovii* Pavlov [Lf: Herb. Lc: Per. D: 28]
 345. *Scutellaria navicularis* Juz. [Lf: Subshrub. Lc: Per. Cs: RB. D: 18, 24]
 346. *Scutellaria subcaespitosa* Pavlov [Lf: Herb. Lc: Per. Cs: RB. D: 27, 28, 29]
 347. *Scutellaria titovii* Juz. [Lf: Herb. Lc: Per. D: 26]
 348. *Scutellaria turgaica* Juz. [Lf: Herb. Lc: Per. D: 9, 10, 10a, 11, 11a, 16]

Gen. 101. *Thymus* L.

349. *Thymus crebrifolius* Klovok [Lf: Dwarf semishrub. Lc: Per. D: 10, 10a, 11a]
 350. *Thymus eremita* Klovok [Lf: Dwarf semishrub. Lc: Per. D: 10a, 11]
 351. *Thymus karatavicus* Dmitrieva [Lf: Dwarf semishrub. Lc: Per. D: 28, 29]
 352. *Thymus magnificus* Klovok [Lf: Dwarf semishrub. Lc: Per. D: 25]

Fam. 21. Liliaceae Juss.

Gen. 102. *Fritillaria* L.

353. *Fritillaria kolbintsevii* Rukšāns & Zubov [Lf: Herb. Lc: Per. D: 24]

Gen. 103. *Gagea* Salisb.

354. *Gagea almaatensis* Levichev, A.Peterson & J.Peterson [Lf: Herb. Lc: Per. D: 25]
 355. *Gagea iliensis* Popov [Lf: Herb. Lc: Per. D: 16, 18]
 356. *Gagea sarysuensis* Murz. [Lf: Herb. Lc: Per. D: 10, 11]
 357. *Gagea ularsaica* I.G.Levichev [Lf: Herb. Lc: Per. D: 28, 29]

Gen. 104. *Tulipa* L.

358. *Tulipa alberti* Regel [Lf: Herb. Lc: Per. Cs: RB. D: 10, 16, 18, 24, 26, 28, 29]
 359. *Tulipa annae* J.de Groot & Zonn. [Lf: Herb. Lc: Per. D: 22, 24]
 360. *Tulipa auliekolica* Perezhogin [Lf: Herb. Lc: Per. D: 2]
 361. *Tulipa berkariensis* Rukšāns * [Lf: Herb. Lc: Per. D: 27, 28, 29]
 362. *Tulipa brachystemon* Regel * [Lf: Herb. Lc: Per. Cs: RB. D: 24]
 363. *Tulipa diana-verettiae* J.de Groot & Zonn. [Lf: Herb. Lc: Per. D: 22]
 364. *Tulipa ivasczenkoae* Epiktetov & Belyalov [Lf: Herb. Lc: Per. D: 24]
 365. *Tulipa kolbintsevii* Zonn. [Lf: Herb. Lc: Per. D: 24]
 366. *Tulipa lemmersii* Zonn., Peterse & J.de Groot [Lf: Herb. Lc: Per. D: 29]
 367. *Tulipa orthopoda* Vved. * [Lf: Herb. Lc: Per. D: 28, 29]
 368. *Tulipa regelii* Krasn. [Lf: Herb. Lc: Per. Cs: RB. D: 26]
 369. *Tulipa turgaica* Perezhogin [Lf: Herb. Lc: Per. D: 9]
 370. *Tulipa salsola* Rukšāns & Zubov [Lf: Herb. Lc: Per. D: 24]

Fam. 22. Nitrariaceae Lindl.

Gen. 105. *Nitraria* L.

371. *Nitraria iliensis* Banaev & Tomoshevich [Lf: Shrub. Lc: Per. D: 18, 24]

Gen. 106. *Tetradiclis* Steven ex M.Bieb.

372. *Tetradiclis corniculata* Khalk. [Lf: Herb. Lc: An. D: 12]

Fam. 23. Orobanchaceae Vent.

Gen. 107. *Euphrasia* L.

373. *Euphrasia integriloba* J.J.Dmitriev & N.I.Rubtzov [Lf: Herb. Lc: An. D: 24]
374. *Euphrasia karataviensis* Govor. [Lf: Herb. Lc: An. D: 28, 29]

Gen. 108. *Pedicularis* L.

375. *Pedicularis czuiliensis* Semiotr. [Lf: Herb. Lc: Per. Cs: RB. D: 26]
376. *Pedicularis interrupta* subsp. *tarbagataica* (Semiotr.) Kamelin [Lf: Herb. Lc: Per. Cs: RB. D: 23]
377. *Pedicularis kokpakensis* Semiotr. [Lf: Herb. Lc: Per. D: 25a]
378. *Pedicularis masalskyi* Semiotr. [Lf: Herb. Lc: Per. D: 29]
379. *Pedicularis transversa* Baimukhambetova [Lf: Herb. Lc: Per. D: 25a]

Fam. 24. Plantaginaceae Juss.

Gen. 109. *Linaria* Mill.

380. *Linaria macrophylla* Kuprian. [Lf: Herb. Lc: Per. D: 8]

Fam. 25. Plumbaginaceae Juss.

Gen. 110. *Acantholimon* Boiss.

380. *Acantholimon karatavicum* Pavlov [Lf: Undershrub. Lc: Per. D: 17, 28]
382. *Acantholimon linczevskii* Pavlov [Lf: Undershrub. Lc: Per. Cs: RB. D: 28]
383. *Acantholimon mikeschinii* Lincz. [Lf: Undershrub. Lc: Per. D: 28]
384. *Acantholimon minshelkense* Pavlov [Lf: Subshrub. Lc: Per. D: 28]
385. *Acantholimon pavlovii* Lincz. [Lf: Undershrub. Lc: Per. D: 29]
386. *Acantholimon squarrosus* Pavlov [Lf: Undershrub. Lc: Per. D: 28]

Gen. 111. *Limonium* Mill.

387. *Limonium botschantzevii* (Lincz.) M.Malekm., Akhani & Borsch [Lf: Herb. Lc: Per. D: 21]
388. *Limonium michelsonii* Lincz. [Lf: Herb. Lc: Per. Cs: RB. D: 24, 25, 25a]

Fam. 26. Poaceae Barnhart

Gen. 112. *Agropyron* Gaertn,

389. *Agropyron cristatum* subsp. *tarbagataicum* (Plotn.) Tzvelev [Lf: Herb. Lc: Per. D: 22, 23]

Gen. 113. *Elymus* L.

390. *Elymus arcuatus* (Golosc.) Tzvelev [Lf: Herb. Lc: Per. D: 25]
391. *Elymus glaucissimus* (Popov) Tzvelev [Lf: Herb. Lc: Per. D: 25]
392. *Elymus sibiricus* Kotukhov [Lf: Herb. Lc: Per. D: 22]

Gen. 114. *Festuca* Tourn. ex L.

393. *Festuca irtyschensis* E.B.Alexeev [Lf: Herb. Lc: Per. D: 3]
394. *Festuca saurica* E.B.Alexeev [Lf: Herb. Lc: Per. D: 23]

Gen. 115. *Leymus* Hochst.

395. *Leymus divaricatus* (Drobow) Tzvelev [Lf: Herb. Lc: Per. D: 25, 28]

Gen. 116. *Limnas* Trin.

396. *Limnas vereschaginii* Krylov & Schischk. [Lf: Herb. Lc: Per. Cs: RB. D: 22]

Gen. 117. *Poa* L.

397. *Poa koksuensis* Golosc. [Lf: Herb. Lc: Per. D: 24]

Gen. 118. *Puccinellia* Parl.

398. *Puccinellia macropus* V.I.Krecz. [Lf: Herb. Lc: Per. D: 26]

Gen. 119. *Stipa* L.

399. *Stipa argillosa* Kotukhov * [Lf: Herb. Lc: Per. D: 22]

400. *Stipa austroaltaica* Kotukhov [Lf: Herb. Lc: Per. Cs: RB. D: 22]

401. *Stipa karakabinica* Kotukhov [Lf: Herb. Lc: Per. D: 23]

402. *Stipa kempirica* Kotukhov [Lf: Herb. Lc: Per. D: 23]

403. *Stipa kotuchovii* M.Nobis [Lf: Herb. Lc: Per. D: 23]

Fam. 27. Polygonaceae Juss.

Gen. 120. *Atraphaxis* L.

404. *Atraphaxis muschketowii* Krasn. [Lf: Shrub. Lc: Per. Cs: RB. D: 25]

405. *Atraphaxis teretifolia* (Popov) Kom. [Lf: Shrub. Lc: Per. Cs: RB. D: 10, 11, 18]

Gen. 121. *Calligonum* L.

406. *Calligonum turbineum* Pavlov [Lf: Shrub. Lc: Per. D: 17, 18, 20]

Gen. 122. *Rumex* L.

407. *Rumex fischeri* Rchb. [Lf: Herb. Lc: Per. D: 24]

408. *Rumex komarovii* Schischk. & Serg. [Lf: Herb. Lc: Per. D: 11]

Fam. 28. Ranunculaceae Juss.

Gen. 123. *Aquilegia* L.

409. *Aquilegia karatavica* Mikeschin [Lf: Herb. Lc: Per. Cs: RB. D: 28]

410. *Aquilegia vitalii* Gamajun. [Lf: Herb. Lc: Per. Cs: RB. D: 24]

Gen. 124. *Delphinium* L.

411. *Delphinium austroaltaicum* A.L.Ebel [Lf: Herb. Lc: Per. D: 22]

412. *Delphinium connectens* Pachom. [Lf: Herb. Lc: Per. D: 25a]

413. *Delphinium pavlovii* Kamelin [Lf: Herb. Lc: Per. D: 28]

Gen. 125. *Ranunculus* L.

414. *Ranunculus karkaralensis* Schegol. [Lf: Herb. Lc: Per. D: 11a]

Gen. 126. *Thalictrum* Tourn. ex L.

415. *Thalictrum bykovii* Kotukhov * [Lf: Herb. Lc: Per. D: 22]

Fam. 29. Rosaceae Juss.

Gen. 127. *Alchemilla* L.

416. *Alchemilla goloskokovii* Juz. [Lf: Herb. Lc: Per. D: 24]

Gen. 128. *Amelanchier* Medik.

417. *Amelanchier turkestanica* Litv. [Lf: Shrub. Lc: Per. D: 11]

Gen. 129. *Cotoneaster* Medik.

418. *Cotoneaster alatavicus* Popov [Lf: Shrub. Lc: Per. D: 24, 25, 27, 29]

419. *Cotoneaster altaicus* G.Klotz ex J.Fryer & B.Hylmö [Lf: Shrub. Lc: Per. D: 25]

420. *Cotoneaster krasnovii* Pojark. [Lf: Shrub. Lc: Per. D: 18, 24, 25, 26]

421. *Cotoneaster neoantoninae* A.N.Vassiljeva [Lf: Shrub. Lc: Per. D: 24, 25]

422. *Cotoneaster polyanthemus* E.L.Wolf [Lf: Shrub. Lc: Per. D: 24, 25]

423. *Cotoneaster talgaricus* Popov [Lf: Shrub. Lc: Per. D: 24, 25, 25a]

Gen. 130. *Crataegus* L.

424. *Crataegus ambigua* subsp. *transcaspica* (Pojark.) K.I.Chr. [Lf: Tree. Lc: Per. D: 13b]

Gen. 131. *Potentilla* L.

425. *Potentilla karatavica* Juz. [Lf: Herb. Lc: Per. D: 28]

426. *Potentilla salsa* Yu.A.Kotukhov [Lf: Herb. Lc: Per. D: 22]

427. *Potentilla schrenkiana* Regel [Lf: Herb. Lc: Per. D: 23, 24]

Gen. 132. *Rosa* L.

428. *Rosa dsharkenti* Chrshan. [Lf: Shrub. Lc: Per. D: 18]

429. *Rosa iliensis* Chrshan. [Lf: Shrub. Lc: Per. D: 17, 18]

430. *Rosa potentilliflora* Chrshan. & Popov [Lf: Shrub. Lc: Per. D: 25]

431. *Rosa schrenkiana* Crép. [Lf: Shrub. Lc: Per. D: 24]

Gen. 133. *Spiraeanthus* (Fisch. & C.A.Mey.) Maxim.

432. *Spiraeanthus schrenkianus* (Fisch. & C.A.Mey.) Maxim. [Lf: Shrub. Lc: Per. Cs: RB. D: 16, 21, 26, 28]

Fam. 30. Rubiaceae Juss.

Gen. 134. *Galium* L.

433. *Galium kasachstanicum* Pachom. [Lf: Herb. Lc: Per. D: 25a]

434. *Galium turgaicum* Knjaz. [Lf: Herb. Lc: Per. D: 7a]

435. *Galium zaisanicum* Pinzhenina & Kupr. * [Lf: Herb. Lc: Per. D: 12]

Gen. 135. *Rubia* L.

436. *Rubia cretacea* Pojark. [Lf: Herb. Lc: Per. Cs: RB. D: 1, 6, 7a, 8, 13]

437. *Rubia pavlovii* Bajtenov & Myrz. [Lf: Herb. Lc: Per. Cs: RB. D: 28]

Fam. 31. Rutaceae Juss.

Gen. 136. *Haplophyllum* A.Juss.

438. *Haplophyllum eugenii-korovinii* Pavlov [Lf: Subshrub. Lc: Per. Cs: RB. D: 28]

439. *Haplophyllum multicaule* Vved. [Lf: Subshrub. Lc: Per. D: 5, 10, 14, 16, 18, 24, 26]

Fam. 32. Scrophulariaceae Juss.

Gen. 137. *Scrophularia* Tourn. ex L.

440. *Scrophularia dshungarica* Golosk. & Tzag. [Lf: Herb. Lc: Per. Cs: RB. D: 24]

441. *Scrophularia nuraniae* Tzag. [Lf: Herb. Lc: Per. Cs: RB. D: 29]

Fam. 33. Thymelaeaceae Juss.

Gen. 138. *Diarthron* Turcz.

442. *Dendrostellera ammodendron* (Kar. & Kir.) Botsch. [Lf: Shrub. Lc: Per. D: 17, 18, 24, 25]

Fam. 34. Zygophyllaceae R.Br.

Gen. 139. *Zygophyllum* L.

443. *Zygophyllum balchaschense* Boriss. [Lf: Herb. Lc: Per. D: 11, 16]

444. *Zygophyllum betpakdalense* Golosk. & Semiotr. [Lf: Herb. Lc: Per. D: 16]

445. *Zygophyllum borissovae* Beier & Thulin [Lf: Herb. Lc: Per. D: 10, 11, 11a, 16]

446. *Zygophyllum furcatum* C.A.Mey. [Lf: Herb. Lc: Per. D: 10, 11, 11a, 23, 24]

447. *Zygophyllum karatavicum* Boriss. [Lf: Herb. Lc: Per. Cs: RB. D: 28]

448. *Zygophyllum kopalense* Boriss. [Lf: Herb. Lc: Per. Cs: RB. D: 16, 18, 25, 26]

449. *Zygophyllum steropterum* Schrenk [Lf: Herb. Lc: Per. D: 16, 17, 18, 26]

450. *Zygophyllum subtrijugum* C.A.Mey. [Lf: Herb. Lc: Per. D: 3, 4, 10, 10a, 11, 16]

451. *Zygophyllum taldykurganicum* Boriss. [Lf: Herb. Lc: Per. D: 18, 24]

Notes*:

- **Arthrophytum subulifolium* Schrenk, according to POWO, is noted in Turkmenistan; however, according to our data and the scientific paper by Osmanali et al. (2019), this indication is incorrect. This species is a narrow-local endemic of the Chu-Ili Range in the south of Kazakhstan (Osmanali et al. 2019).
- **Atriplex iljinii* Aellen, similar to *Arthrophytum subulifolium* Schrenk, is erroneously reported in Turkmenistan, as per the POWO database. The distribution of *A. iljinii* is poorly studied, known mainly from type specimens. This species is observed in the northwest of the Mangistau Region, as well as in the Aktobe and Kyzylorda (Aralkum Desert) Regions of Kazakhstan (Suchorukow 2007).
- **Ferula leucographa* Korovin, according to POWO, is recorded in Uzbekistan; yet, according to the updated synopsis of *Apiaceae* of Kazakhstan and Central Asia (Pimenov 2020), the species grows only in Kazakhstan.
- **Rindera ochroleuca* Kar. & Kir., as suggested by POWO, is observed in Altai in the Russian Federation; yet, our investigation did not find credible evidence supporting this information.
- **Eutrema halophilum* (C.A.Mey.) Al-Shehbaz & Warwick was previously reported in China (Wu et al. 2008); however, German DA and Chen WL (2009) in their scientific paper did not confirm the presence of this species in China.
- **Astragalus leucocalyx* Popov is recorded in POWO for Uzbekistan; nevertheless, Tojibaev et al. (2020b) in their scientific paper exclusively listed this species as occurring in Kazakhstan.
- **Prangos multicostata* Kljuykov & Lyskov., according to the scientific paper by Pimenov (2020), is a synonym for *Prangos dzhungarica* Pimenov. Further study is required to investigate whether species status of *Prangos multicostata* is warranted.
- **Centaurea kryloviana* Serg. is not recognised in POWO and GBIF, probably due to nomenclature errors. However, according to the scientific paper by Kupriyanov (2018), this species is endemic to eastern Kazakhstan.
- **Astragalus rariflorus* Ledeb., previously noted for Western Siberia (Krylov 1933), yet we did not find herbarium materials of this species from this territory. This species is also not recorded in the Flora of Siberia (Malyshev 1994).
- **Lappula cristata* (Bunge) B. Fedtsch. in JBIF is recorded for Western Siberia, on the territory of the Russian Federation (Bochkov and Seregin 2022); however, the presence of this species lacks substantial verification. Notably, the Flora of Siberia (Malyshev 1997) does not include any records of this species. Therefore, further investigation is necessary to elucidate the occurrence of *Lappula cristata* within the territory of the Russian Federation.
- **Onobrychis alata* Bajtenov was omitted from the list of flora of Kazakhstan (Abdulina 1999). *Conspectus Florae Asiae Mediae* (Kamelin et al. 1981) noted that this species needs to be re-collected to confirm species status.
- **Tulipa berkariensis* Rukšāns in POWO is recognised as a synonym of *Tulipa kaufmanniana* Regel., based on data from Everett (2013). Nevertheless, *T. berkariensis* from the Berkara Valley and other places in Kazakhstan has a lower amount of nuclear 2C DNA (based on the data flow cytometric measurement of nuclear DNA content) than authentic *T. kaufmanniana* from Uzbekistan (Zonneveld 2009). This discrepancy implies that *Tulipa berkariensis* should be recognised as a distinct taxonomic entity.

- **Tulipa orthopoda* Vved. listed in POWO is recognised as a synonym for *Tulipa bifloriformis* Vved., based on Christenhusz et al. (2013) and Everett (2013). This scientific paper also notes that the species status of *Tulipa orthopoda* should be warranted, based on differences in morphological characters and flowering period, but further fieldwork is required to establish the variability of *T. bifloriformis* in the wild.
- **Tulipa brachystemon* Regel in POWO is recognised as a synonym for *Tulipa tetraphylla* Regel, also according to Christenhusz et al. (2013) and Everett (2013). However, Zonneveld (2009) distinguishes *Tulipa brachystemon* Regel as an independent taxon, based on the nuclear DNA content (DNA value 2C).
- **Galium zaisanicum* Pinzhenina & Kupr. was described quite recently (Pinzhenina and Kupriyanov 2023), presumably due to its recent identification it is absent in the GBIF and POWO systems.
- **Stipa argillosa* Kotukhov in POWO is erroneously listed for the Altai Republic of the Russian Federation. This species is described from the territory of East Kazakhstan. Type: Southern Altai, south-eastern foothills of the Azutau Ridge, Bulgartabat tract, foothill desert, outcrops of tertiary clays, clay-rub-bly areas, 05/22/1991, Yu. Kotukhov (LE) (Kotukhov 1998). The Azutau Ridge borders the basin of Lake Markakol from the south and is entirely situated within the territory of Kazakhstan (Yegorina et al. 2003).
- **Thalictrum bykovii* Kotukhov, as well as the previous species in POWO, is erroneously listed for the Altai Republic of the Russian Federation. This species was also described from the territory of East Kazakhstan. Type: Southern Altai, eastern spurs of Azutau Ridge, Mramornaya Mount, Middle belt, 900–1100 m above sea level, south-eastern slope, steppe shrub meadows, 14/06/1984, Yu. Kotukhov (LE).

Supplementary material 1

Checklist of subendemic taxa previously considered endemic in Kazakhstan

Authors: Serik A. Kubentayev, Daniyar T. Alibekov, Yuri V. Perezhogin, Georgy A. Lazkov, Andrey N. Kupriyanov, Alexander L. Ebel, Klara S. Izbastina, Olga V. Borodulina, Balsulu B. Kubentayeva

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.238.114475.suppl1>

Supplementary material 2

Former endemics of Kazakhstan that are now reclassified as synonyms for species exhibiting broader geographical distributions

Authors: Serik A. Kubentayev, Daniyar T. Alibekov, Yuri V. Perezhogin, Georgy A. Lazkov, Andrey N. Kupriyanov, Alexander L. Ebel, Klara S. Izbastina, Olga V. Borodulina, Balsulu B. Kubentayeva

Data type: docx

Copyright notice: This dataset is made available under the Open Database License (<http://opendatacommons.org/licenses/odbl/1.0/>). The Open Database License (ODbL) is a license agreement intended to allow users to freely share, modify, and use this Dataset while maintaining this same freedom for others, provided that the original source and author(s) are credited.

Link: <https://doi.org/10.3897/phytokeys.238.114475.suppl2>